



PS Clethodim

| ACTIVE INGREDIENT: | % by Weight |
|--------------------------|-------------|
| *Clethodim | 26.4% |
| OTHER INGREDIENTS: | 73.6% |
| TOTAL: | 100.0% |

Contains Petroleum Distillates

*(E)-2-[1[[[(3-chloro-2-propenyl)-oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one

Contains 2.0 lbs. clethodim per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for additional PRECAUTIONARY STATEMENTS.

EPA Reg. No. 89442-4

Net Contents: 1 Gallon

20121210

Manufactured for:
Prime Source, LLC

4609 E. Boonville-New Harmony Road
Evansville, IN 47725

EPA Est. No. 70989-MO-001

| FIRST AID | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person. |
| IF INHALED: | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| HOT LINE NUMBER | |
| <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222.</p> | |
| NOTE TO PHYSICIAN | |
| <p>Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. If ingested, probable mucosal damage may contraindicate the use of gastric lavage.</p> | |

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

Causes substantial but temporary eye irritation. Avoid contact with skin. Do not get in eyes, on skin or on clothing. Harmful if swallowed or inhaled. Avoid breathing vapors or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves such as Barrier Laminate or Viton \geq 14 mils, shoes plus socks, and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwater or rinsate.

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist:

| | |
|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Solano Grass: | Solano County, California: the vernal lakes are bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south, and Travis Air Force Base to the west. |
| Wild Rice: | Hays County, Texas. |

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves, such as Barrier Laminate or Viton \geq 14 mils, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift. Do not enter treated areas without protective clothing until sprays have dried.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank with this product.

CHEMIGATION

Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed. Refer to supplemental labeling entitled, "Application of PS Clethodim Onions (dry bulbs and green) and Garlic by Chemigation", for use directions for chemigation.

May be applied to onions and garlic by sprinkler irrigation systems. Do not apply by chemigation to any other crop, or to this crop using any other type of irrigation system.

PRODUCT INFORMATION

FOR USE ON: Alfalfa, Artichoke (Globe), Asparagus, Bean and Pea (dry shelled)¹, Bean and Pea (succulent)², Broccoli, Bushberry³, Cabbage, Caneberry⁴, Canola⁴, Carrot, Cauliflower (and other Head and Stem Brassica Vegetables)⁵, Celery, Clover (grown in Idaho, Oregon and Washington only), Conifers, Cotton, Cranberry, Cucumber, Eggplant (and other Fruiting Vegetables)⁶, Fallow Land (and other non-producing agricultural areas), Flax⁴, Field Corn⁷, Garden Beet, Garlic, Herbs⁸, Hops, Horseradish (and other Root Vegetables)⁹, Legume Vegetables (edible podded)¹⁰, Lettuce, Head and Leaf (and other leafy greens)¹¹, Melons (including Cantaloupes and Watermelons)¹², Mint, Mustard Greens (and other leafy brassica greens)¹³, Mustard Seed⁹, Non-Bearing Food Crops, Non-Crop or Non-Planted Areas, Onions (dry bulb and green), Ornamentals, Peaches, Peanut (including perennial), Peppers (bell and non-bell), Potato, Radish, Rhubarb (and other Leaf Petioles)¹⁴, Safflower, Sesame, Shallots (dry bulbs and green), Soybeans, Squash (including Pumpkins), Strawberry, Sugar Beet, Sunflower, Sweet Potato, Tomato, Turnip Greens, Yam (and other Tuberous and Corn Vegetables)¹⁵

*Not for use in California

- ¹ Other Dry Shelled Bean and Pea crops approved for use with PS Clethodim include: Bean (*Lupinus* spp.), grain, sweet, white and white sweet; Bean (*Phaseolus* spp.), field, kidney, lima (dry), navy, pinto and tepary; Bean (*Vigna* spp.), adzuki bean, black-eyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, broad (dry), chickpea (garbanzo), guar, lablab bean and lentil; Pea (*Pisum* spp.), field and pigeon.
- ² Other Succulent Bean and Pea crops approved for use with PS Clethodim include: Bean (*Lupinus* spp.), grain, sweet, white and white sweet; Bean (*Phaseolus* spp.), field, kidney, lima (dry), navy, pinto and tepary; Bean (*Vigna* spp.), adzuki bean, black-eyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, broad (dry), chickpea (garbanzo), guar, lablab bean and lentil; Pea (*Pisum* spp.), field and pigeon.
- ³ Other bushberry crops approved for use with PS Clethodim include: Aronia Berry, Blueberry (highbush), Black Currant, Buffalo Currant, Chilean Guava, Cranberry (highbush), Elderberry, European Barberry, Gooseberry, Honeysuckle (edible), Huckleberry (edible), Jostaberry, Juneberry (Saskatoon berry), Native Currant, Salal, Sea Buckthorn, Red Currant; cultivars, varieties and/or hybrids of these.
- ⁴ Other caneberry crops approved for use with PS Clethodim include: Blackberry, Loganberry, Raspberry (black and red), Wild Raspberry; cultivars, varieties and/or hybrids of these.
- ⁵ Other head and stem brassica vegetables approved for use with PS Clethodim include: Chinese broccoli, Brussels sprouts, Chinese (napa) cabbage, Chinese mustard, cavalo broccoli and kohlrabi.
- ⁶ Other Fruiting Vegetables (except tomato) approved for use with PS Clethodim include: eggplant, groundcherry, pepino, peppers (all) and tomatillo.
- ⁷ For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn.
- ⁸ Other Herb crops approved for use with PS Clethodim include: angelica, balm, basil, borage, burnet, chamomile, catnip, chervil (dried), chive, Chinese chive, clary, coriander (leaf), costmary, cilantro (leaf), curry (leaf), dill (dillweed), horehound, hyssop, lavender, lovage (leaf), marigold, marjoram (*origanum* spp.), nasturtium, parsley (dried), pennyroyal, rosemary, rue, sage and savory, summer and winter.
- ⁹ Other root vegetables approved for use with PS Clethodim include: burdock, edible; celeriac; chervil, turnip-rooted; chycory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify, black; salsify, Spanish; skirret and turnip.
- ¹⁰ Other Edible Podded Legume Vegetable crops approved for use with PS Clethodim include: Bean (*Phaseolus* spp.), runner, snap and wax; Bean (*Vigna* spp.), asparagus, Chinese longbean, moth, yardlong, jackbean; Pea (*Pisum* spp.), dwarf, edible-pod, snow, sugar snap, pigeon and sword bean.
- ¹¹ Other Leafy Greens crops approved for use with PS Clethodim include: amaranth (Chinese spinach, leafy amaranth and tampala), arugula (rockette), chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden, yellow rock

- and winter), dandelion, dock (sorrel), endive (escarole), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio (red chicory), spinach (New Zealand and Vine (Indian and Malabar).
- ¹² Other cucurbit crops approved for use with PS Clethodim include: Chayote (fruit), Chinese Wax Gourd, Citron Melon, Edible Gourd, Gherkin and Muskmelons (all) including Honeydew Melon.
- ¹³ Other leafy brassica greens approved for use with PS Clethodim include: broccoli raab, Chinese (bok choy) cabbage, collards, kale, mizuna, mustard greens, mustard spinach and rape greens.
- ¹⁴ Other leaf petiole crops approved for use with PS Clethodim include: cardoon, celtuce, Chinese celery, Florence fennel, and Swiss chard.
- ¹⁵ Other tuber and corm vegetables approved for use with PS Clethodim include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible burdock, edible canna, bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, taniar, turmeric and bean yam.

PS Clethodim is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

PS Clethodim is a selective postemergence herbicide for control of annual and perennial grasses. PS Clethodim does not control sedges or broadleaf weeds.

Repeated use of PS Clethodim (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species.

If poor performance occurs and cannot be attributed to adverse weather or application conditions, a resistant biotype may be present. This is most likely to occur in fields where other control strategies such as crop rotation, mechanical removal, and other classes of herbicides are not used from year to year.

Do not allow PS Clethodim to come in contact with desirable grass crops such as corn, rice, sorghum, small grains, or turf, as these and other grass crops will be injured or killed. Minor leaf spotting may occur on treated plants under certain environmental conditions. New foliage is not affected.

Control Symptoms

Treated grass weeds show a reduction in vigor and growth. Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days after application, depending on grass species treated and environmental conditions.

APPLICATION INFORMATION

Timing of Applications

Apply PS Clethodim postemergence to actively growing grasses according to rate table instructions. Applications made to grass plants stressed by insufficient moisture, or hot or cold temperatures, or to grass plants exceeding listed growth stages may result in unsatisfactory control. Do not apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, PS Clethodim should be applied as soon as possible, after irrigation (within 7 days). In arid regions, a second application of PS Clethodim will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of PS Clethodim may reduce weed control. DO NOT APPLY PS Clethodim if rainfall is expected within one hour, since control may be reduced.

ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE

| CROP | ADJUVANT USE INSTRUCTIONS |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Alfalfa, Cotton, Dry Shelled Bean & Pea, Edible Poddled Legume Vegetable, Peanuts (including perennial), Potato, Soybean, Succulent Bean & Pea, Sugar Beet and Sunflower | Always use a crop oil concentrate* at 1.0 qt./A by ground or 1% v/v (but not less than 1 pt./A) in the finished spray volume by air. 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to PS Clethodim applications, in addition to the recommended rate of crop oil concentrate. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |
| Artichoke (Globe), Asparagus, Bushberry, Caneberry, Canola, Carrot, Clover, Cranberry, Cucurbits, Flax, Fruiting Vegetables (except Tomato), Garden Beet, Garlic, Head & Stem Brassica Vegetables, Herbs, Hops, Leaf Petioles, Leafy Brassica Greens, Leafy Greens, Mint, Mustard Seed, Onion (Dry Bulb & Green), Peaches, Root Vegetables, Safflower, Sesame, Shallots (Dry Bulb & Green), Strawberry, Sweet Potato (Yam & other Tuberous and Corm Vegetables except Potato) and Tomato | Always use a crop oil concentrate at 1% v/v in the finished spray volume unless tank mix instructions indicate otherwise. Addition of liquid fertilizer is not recommended for these crops. |
| Non-Bearing Food Crops, Ornamental Plants | Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v). Use of crop oil concentrate is not recommended since it may injure flowers and foliage. |
| Conifer Trees, Fallow Land (and other non-producing agricultural areas), and Non-Crop or Non-Planted Areas | Always use a crop oil concentrate containing at least 15% emulsifier at 1% v/v (but not less than 1 pt./A) in the finished spray volume. |

*Acceptable crop oil concentrates would be those that contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality, and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

Ground Application

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 40 gals. of spray solution per acre. Under the following conditions a minimum of 10 gals. per acre is required: ultra narrow row cotton, narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions,

heavy grass pressure or when grasses are at or near maximum height. Failure to use a minimum of 10 gals. per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Applications to onions (dry bulbs and green), garlic, and shallots (dry bulbs and green) should be made in a minimum of 20 gals. of spray solution per acre.

Air Application

Use a minimum of 3 gals. of spray solution per acre unless otherwise directed in this label. Increase spray volumes up to 10 gals. as grass or crop foliage becomes dense. For onions (dry bulbs and green), garlic, or shallots (dry bulbs and green). When applying by air do not exceed 8 fl. oz./A in a single application. In California, air applications to onions, garlic or shallots should be made in a minimum of 20 gals. of spray solution per acre. In states other than California, air application to onions, garlic, or shallots should be made in a minimum of 10 gals. of spray solution per acre.

NOTE: Crop injury may occur when PS Clethodim is applied to onions, garlic or shallots with aerial equipment.

Spot Treatment

When using hand sprayers or high volume sprayers utilizing hand guns, mix 1/4% to 1/2% (0.33 oz. to 0.65 oz. per gal.) PS Clethodim and treat to wet vegetation, while not allowing runoff of spray solution. For uses requiring crop oil concentrate, include crop oil concentrate at 1% (1.3 oz. per gal.) by volume. For uses requiring non-ionic surfactant, include non-ionic surfactant at 1.4% (0.33 oz. per gal.) by volume).

NOTE: If PS Clethodim is applied as a spot treatment, care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.

CHEMIGATION – ONION (Dry Bulb and Green) AND GARLIC SPRINKLER IRRIGATION APPLICATION

Do not apply PS Clethodim by chemigation in the states of Idaho, Montana, Oregon and Washington

Apply PS Clethodim at the high rate recommended for annual grasses (16 fl. oz. per acre) when the grass height is at the low end of the range (application to larger grasses may not provide adequate control). Add a crop concentrate containing at least 15% emulsifier at 1 quart per acre.

Apply PS Clethodim in 0.1 to 0.2 acre-inch of water either at the end of a regular irrigation set or as a separate application not associated with a regular irrigation using the least amount of water that provides proper distribution and coverage. Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness. Use a metering device to inject the PS Clethodim into the irrigation water at a constant flow. Constant agitation must be maintained in the chemical supply tank during the entire period of herbicide application. Inject the product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period.

It is not recommended that PS Clethodim be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Use Precautions

1. Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
3. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.
5. A person knowledgeable of the chemigation system and responsible for its operation or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
12. Do not apply when wind speed favors drift beyond the area intended for treatment.

RESTRICTIONS AND LIMITATIONS

Do not apply if rain is expected within 1 hour of application as control may be unsatisfactory.

Do not apply a postemergence broadleaf herbicide within one day following application of PS Clethodim or reduced grass control may result.

PS Clethodim is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

For canola, do not apply more than 6 fl. oz. of PS Clethodim per acre per season. For clover, flax, mustard seed and radish crops, do not apply more than 16 fl. oz. of PS Clethodim (0.25 lb. ai) per acre per season. For all other crops, do not apply more than 32 fl. oz. of PS Clethodim (0.50 lb. ai) per acre per season. Application on Long Island, New York, is restricted to no more than 16 fl. oz. of PS Clethodim (0.25 lb. ai) per acre per season.

Do not apply more than 8 fl. oz./A of PS Clethodim per application to the following crops: asparagus, brassica vegetables (head

and stem), bean (succulent), carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beet, green onion, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, legume vegetables (edible podded), non-bearing food crops, pea (dry shelled), pea (succulent), root vegetables, safflower, sesame and strawberry. Do not apply more than 6 fl. oz./A of PS Clethodim per application to canola or mustard seed. For all other crops, do not apply more than 16 fl. oz. of PS Clethodim (0.25 lb. a) per acre per application. **Exceeding these rates may result in unacceptable crop injury.**

Do not apply under conditions of stress. Applying PS Clethodim under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate PS Clethodim effectively, and will be less susceptible to herbicide activity.

Optimal perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices, (discing, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, such as continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, result in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, no fewer than two PS Clethodim applications per season per year are recommended at the appropriate weed-growth stage rate under continuous no-till conditions.

Grass crops such as corn, rice, sorghum, small grains, or turf, etc. are highly sensitive to PS Clethodim.

While all the vegetable crops on this label have been tested and are tolerant to PS Clethodim, not all specialty varieties of these crops have been tested. It is advised that, before applying PS Clethodim to specialty varieties of vegetable crops on this label, crop tolerance be investigated first using a small section of the field. It is possible that injury symptoms can occur. Symptoms may appear as leaf speckling or stunting.

Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.

Tank mixes of PS Clethodim and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of PS Clethodim may be necessary. **AVOID SPRAY DRIFT.**

Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:

- Do not spray if wind speeds are or become excessive.
- Do not spray if wind speed is 10 mph or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions.
- Do not spray if winds are gusty.
- Use extreme caution when conditions are favorable for drift (high temperatures, drought, low relative humidity), especially when sensitive plants are located nearby.
- Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.
- Further reductions in drift can be obtained by:
 1. Using large droplet size sprays. Do not use nozzles that produce small droplets. Orient nozzles downward and slightly backward as needed to reduce drift for ground applications.
 2. Orienting nozzles straight back with windstream, using straight stream orifices for aerial applications. Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 3 GPA volume. Application height and boom length should be set according to manufacturer's instructions to minimize drift.
 3. Increasing the volume of spray mixture (for example, a minimum of 10 GPA for ground applications) by using higher flow

rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift.

4. Applying as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR PS CLETHODIM

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|----------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Alfalfa including: Sainfoin Holy Clover Birdsfoot trefoil ⁽³⁾ | 15 days before grazing, feeding or harvesting (cutting) for forage or hay | 6-16 fl. oz ⁽⁴⁾ | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air ⁽⁵⁾ | Refer to tank mix partners for feeding, grazing and harvesting restrictions. ^(5,6) The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |
| Artichoke (Globe) | 5 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. Do not apply more than 32 fl. oz./A (0.5 lb. ai/A) per season. |
| Asparagus | 1 day | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |

| Crops ⁽¹⁾ | Minimum Time From Application to Harvest (PHI) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre ⁽²⁾ | Special Use Instructions |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------|--------------------|------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>Bean, Dry Shelled including: Bean (<i>Lupinus</i> spp.) Grain, Sweet, White, White Sweet Bean (<i>Phaseolus</i> spp.) Field, Kidney, Lima (dry), Navy, Pinto, Tepary Bean (<i>Vigna</i> spp.) Adzuki Bean Black-eyed Pea Catjang, Cowpea, Crowder Pea Moth Bean Mung Bean Rice Bean Southern Pea Urd Bean Broad (dry) Chickpea (garbanzo) Guar Lablab Bean Lentil</p> | 30 days | 6-16 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | <p>Do not apply more than 16 fl. oz./A per application. For repeat applications make on a minimum of 14 day interval. Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn.</p> |

| Crops ⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre ⁽²⁾ | Special Use Instructions |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------|--------------------|------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Bean, Succulent including: Bean (<i>Phaseolus</i> spp.) Broad Bean (succulent) Lima Bean (green) Bean (<i>Vigna</i> spp.) Black-eyed Pea Cowpea Southern Pea | 21 days | 6-8 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | Refer to appropriate Table for reduced rate directions for the control of small annual grasses. Do not apply more than one (1) application per acre per season. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn. |
| Beet, Garden | 30 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Brassica Vegetables, Head & Stem including: Broccoli Brussels sprouts Cabbage Cauliflower | 30 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |

| Crops ⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre ⁽²⁾ | Special Use Instructions |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------|--------------------|----------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Bushberry, including: Aronia berry Blueberry, highbush Chilean Guava Cranberry, Highbush Currant, Black Currant, Buffalo Currant, native Currant, Red Elderberry European Barberry Gooseberry Honeysuckle, edible Huckleberry Jostaberry Juneberry Saskatoon berry Native currant Salal Sea Buckthorn Cultivars, varieties and/or hybrids of these. | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. Do not apply more than 32 fl. oz. (0.5 lb. ai/A) per season. For repeat applications make on a minimum of a 14 day interval. Apply at the base of the plant where grassy weeds are growing near the ground. Do not apply to low growing berries. Do not apply to Bushberry grown for root stock. |
| Caneberry, including: Blackberry Loganberry Raspberry, black Raspberry, red Raspberry, wild Cultivars, varieties and/or hybrids of these. | 7 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. Do not apply more than 32 fl. oz./A (0.5 lb. ai/A) per season. For repeat applications make on a minimum of a 14 day interval. Apply at the base of the plant where grassy weeds are growing near the ground. Do not apply to low growing berries. Do not apply to Caneberry grown for root stock. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|---------------------------|--------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Canola | 70 days | 4-6 fl. oz. | 1% v/v in the finished spray volume. | Do not apply after crop has begun bolting. Crop injury may occur when PS Clethodim is applied during the bloom period. Do not apply more than 6 fl. oz./A in a single application. Do not exceed 6 fl. oz/A in a season. |
| Carrot | 30 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Clover | 15 days before grazing, feeding, or harvesting (cutting) for forage or hay | 6-16 fl. oz. | 1% v/v in the finished spray volume. | For use on clover grown in the states of Idaho, Oregon and Washington only. Do not exceed 16 fl. oz/A in a season. |
| Corn, Field For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn. See Directions for Use in Roundup Ready Field Corn (Burndown) table. | 90 days | 3 fl. oz. | Do not use COC or MSO with PS Clethodim in this use pattern. | Do not make more than 1 application per season. Do not apply more than 3 fl. oz./A per season. To control the existing stand, replant no sooner than 6 days after application. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PHI) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|---------------------------|----------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cotton | 60 days | 6-16 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | Do not graze treated fields or feed treated forage or hay to livestock. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |
| Cranberry | 30 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. Do not apply between the "hook" stage and full fruit set. For repeat applications make on a minimum of a 14 day interval. |
| Cucurbits including: Chayote (fruit) Chinese Wax Gourd Citron Melon Cucumber Gherkin Gourd, Edible Muskmelons (all) including: Cantaloupes Honeydew Melon Pumpkin Squash (all) Watermelon | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Fallow Land Conifer Trees (and other non-producing agricultural areas) Non-Crop or Non-Planted Areas | N/A | 6-16 fl. oz. | 1% v/v (but not less than 1 pt./A) in the finished spray volume using a crop oil concentrate containing at least 15% emulsifier. | Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PHI) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|--------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|---------------------------|----------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Flax | 60 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Apply prior to bloom. Crop injury may occur when PS Clethodim is applied during the bloom period. Do not apply more than 8 fl. oz./A in a single application. Do not exceed 16 fl. oz./A in a season. |
| Fruiting Vegetable (except Tomato) including: Eggplant Groundcherry Pepino Peppers (all) Tomatillo | 20 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------|---------------------------|----------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Herbs including: Angelica, Balm, Basil, Borage, Burnet, Camomile, Catnip, Chervil (dried), Chive, Chinese Clary, Coriander (leaf), Costmary, Cilantro (leaf), Curry (leaf), Dill (dillweed), Horehound, Hyssop, Lavender, Lovage (leaf), Marigold, Marjoram (<i>Origanum</i> spp.), Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory, Summer and Winter | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | PS Clethodim has not been tested on all herbs, and herb varieties. It is the responsibility of the user to test PS Clethodim on a small portion of the crop to be treated before treating the entire field. Crop tolerance should be verified to PS Clethodim on a small area of the herb crop, at the desired PS Clethodim rate and with the same crop oil concentrate that will be used on the herb field. If no crop response is evident seven (7) days after treatment, PS Clethodim may be used on the entire field at the rate tested and with the same crop oil used in the tolerance test. Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Hops | 21 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PHI) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|---------------------------|----------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------|
| Leaf Petioles including: Cardoon Celery Celtuce Chinese Celery Fennel, Florence (finocchio) Rhubarb Swiss Chard | 30 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Leafy Brassica Greens including: Broccoli Raab Cabbage, Chinese (bok choy) Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in the single application. For repeat applications make on a minimum of a 14 day interval. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PHI) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|---------------------------|----------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|
| Leafy Greens including: Amaranth Chinese Spinach Leafy Amaranth Tampala Arugula (roquette) Chervil Chrysanthemum, Edible-leaved Chrysanthemum, Garland Corn Salad Cress Garden Upland (yellow rock and winter) Dandelion Dock (sorrel) Endive (escarole) Lettuce, Head and Leaf Orach Parsley Purslane Garden Winter Radicchio (red chicory) Spinach New Zealand Vine (Indian and malabar) | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |

| Crops ⁽¹⁾ | Minimum Time From Application to Harvest (PHI) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre ⁽²⁾ | Special Use Instructions |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------|-----------------------------|------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>Legume Vegetables, Edible Poddeds including: Bean (<i>Phaseolus</i> spp.) Runner Snap Wax Bean (<i>Vigna</i> spp.) Asparagus Chinese Longbean Moth Yardlong Jackbean Pea (<i>Pisum</i> spp.) Dwarf Edible-pod Snow Sugar Snap Pigeon Sword Bean</p> | 21 days | 6-8 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | <p>Do not apply more than 8 fl. oz./A in a single application. Do not apply more than one (1) application per acre per season. For peas apply before bloom, but no later than 21 days before harvest. Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> |
| Mint | 21 days | 6-16 fl. oz. ⁽⁴⁾ | 1 qt. by ground or 1% v/v (but not less than 1 pt./A by air.) | <p>Do not apply more than 16 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval.</p> |
| Mustard Seed | 75 days | 4-6 fl. oz. | 1% v/v in the finished spray volume. | <p>Do not apply after crop has begun bolting. Crop injury may occur when PS Clethodim is applied during the bloom period. Do not apply more than 6 fl. oz./A in a single application. Do not exceed 16 fl. oz./A in a season.</p> |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|---------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------|----------------------------------|--------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Onion (Dry Bulb Only) Garlic Shallot (Dry Bulb Only) | 45 days | 6-16 fl. oz. ^{(7), (8)} | 1% v/v in the finished spray volume. | Minimum of 20 gals./A spray volume by ground in entire U.S. Minimum of 20 gals./A spray volume by air in California. ⁽⁹⁾ In states other than California, air application to onions, garlic or shallots should be made in a minimum of 10 gals./A. |
| Onion, Green including: Green Eschalot Japanese Bunching Onion Leeks Scallion or Spring Onion | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Ornamentals | N/A | 6-16 fl. oz. | Use of crop oil concentrate is not recommended since it may injure flowers and foliage. See Special Use Instructions. | Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v). Sugar maples cannot be tapped for syrup within one year of PS Clethodim application. |
| Non-Bearing Food Crops | N/A | 6-8 fl. oz. ⁽⁸⁾ | See Special Use Instructions. | Do not apply more than 8 fl. oz./A in a single application to non-bearing food crops. |

| Crops ⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre ⁽²⁾ | Special Use Instructions |
|-------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------|--------------------|------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pea, Dry Shelled including: Pea (<i>Pisum</i> spp.) Field Pigeon | 30 days | 6-8 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | Do not apply more than 8 fl. oz./A per application. Do not apply more than one (1) application per acre per season. Apply before bloom but not later than 30 days prior to harvest. ⁽¹⁰⁾ Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn. |
| Pea, Succulent including: Pea (<i>Pisum</i> spp.) English Pea Garden Pea Green Pea Pigeon Pea | 21 days | 6-8 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | Do not apply more than 8 fl. oz./A per application. Do not apply more than one (1) application per acre per season. Apply before bloom but not later than 21 days prior to harvest. ⁽¹⁰⁾ Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|-----------------------------------------------------------------------------------------------------|------------------------------------------------------|---------------------------|------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Peach | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. Do not apply more than 32 fl. oz./A (0.5 lb. ai/A) per season. For repeat applications make on a minimum of a 14 day interval. Do not apply to Peach grown for root stock. |
| Peanut (including Perennial) | 40 days | 6-16 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |
| Potato | 30 days | 8-16 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |
| Radish | 15 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. Do not apply more than 16 fl. oz./A (0.25 lb. ai) per acre in a season. For repeat applications make on a minimum of a 14 day interval. |
| Root Vegetable (except Radish), including: Chicory, Ginseng, Horseradish, Turnip | 30 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |

| Crops⁽¹⁾ | Minimum Time From Application to Harvest (PH) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre⁽²⁾ | Special Use Instructions |
|----------------------------|------------------------------------------------------|---------------------------|------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Safflower | 70 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Sesame | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply during flowering. Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Soybean | 60 days | 6-16 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | Do not graze treated fields or feed treated forage or hay to livestock. Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |
| Strawberry | 4 days | 6-8 fl. oz. | 1% v/v in the finished spray volume. | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14 day interval. |
| Sugar Beet | 40 days | 6-16 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽⁵⁾ | Refer to appropriate Table for reduced rate directions for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |

| Crops ⁽¹⁾ | Minimum Time From Application to Harvest (PHI) | Use Rates Per Acre | Crop Oil Concentrate Rates Per Acre ⁽²⁾ | Special Use Instructions |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------|--------------------|------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sunflower | 70 days | 6-16 fl. oz. | 1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽³⁾ | The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. |
| Sweet Potato, Yam and Other Tuberous and Corn Vegetables (except Potato) including: Artichoke Chinese, Jerusalem Cassava Bitter, Sweet Ginger | 30 days | 6-16 fl. oz. | 1% v/v in the finished spray volume. | The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn. |
| Tomato | 20 days | 6-16 fl. oz. | 1% v/v in the finished spray volume. | For repeat applications make on a minimum of a 14 day interval. |
| Turnip Greens | 14 days | 6-8 fl. oz. | 1% v/v in the finished spray volume | Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval. |

N/A = Not Applicable

- ¹ PS Clethodim is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.
- ² Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the Addition of Adjuvant and Crop Oil Concentrate section for further information.
- ³ PS Clethodim may be applied to seedling or established alfalfa grown for seed, hay, silage, green crop or direct grazing.
- ⁴ For weed control in established alfalfa and mint, the minimum use rate is 10 fl. oz./A.
- ⁵ 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to PS Clethodim applications, in addition to the recommended rate of crop oil

- concentrate.
- ⁶ Do not apply PS Clethodim plus 2,4-DB as a tank mix to alfalfa unless the 60 day feeding, grazing, and harvesting restriction on the 2,4-DB label can be observed.
 - ⁷ For ground applications to garlic or shallot, do not exceed 8 fl. oz./A in a single application. For air applications to onion, garlic or shallot, do not exceed 8 fl. oz./A in a single application. For garlic and shallot, do not exceed 2 applications per season. In CA for air applications to onion, do not exceed 2 applications per season.
 - ⁸ If PS Clethodim is applied as a spot treatment to onion, garlic, shallot, or non-bearing food crops, care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.
 - ⁹ In California, do not apply PS Clethodim to onion, garlic, or shallot until the crop has at least two full leaves. In California, 14 day spray intervals are recommended between the application of PS Clethodim and liquid nitrogen or other herbicide applications. Injury to crop may occur when shorter intervals are observed.
 - ¹⁰ Applications of PS Clethodim to peas during the bloom period could result in severe crop injury, including loss of yield and delayed maturity.

IMPORTANT

Plant tolerance to PS Clethodim at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if the herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of PS Clethodim have investigated the safety factor to plants not listed on the label.

NON-BEARING FOOD CROPS

PS CLETHODIM SHOULD NOT BE APPLIED TO NON-BEARING FRUIT OR NUT CROPS WHICH ARE GROWN FOR ROOT STOCK.

Crop injury to non-bearing fruit and nut crops can occur if PS Clethodim is improperly applied. PS Clethodim should not be applied directly over the top of these plant types. Instead, spray should be directed at the base of the plant where grassy weeds are growing near the ground.

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following PS Clethodim application.

| COMMON NAME | SCIENTIFIC NAME |
|------------------|--------------------------------------------|
| Apples | <i>Malus</i> spp. |
| Berries | <i>Vaccinium</i> spp. <i>Rubus</i> spp. |
| Cherry, Sweet | <i>Prunus avium</i> |
| Citrus Fruits | <i>Citrus</i> spp. |
| Grapes | <i>Vitis</i> spp. |
| Olives | <i>Olea</i> spp. |
| Peach | <i>Prunus persica</i> |
| Pears | <i>Pyrus communis</i> |
| Prunes | <i>Prunus</i> spp. |
| Stone Fruits | <i>Prunus</i> spp. |
| Strawberries | <i>Fragaria</i> spp. |
| Tree Nuts | |
| Almond | <i>Prunus triloba</i> |
| Filbert | <i>Corylus maxima</i> |
| Pecan | <i>Carya illinoensis</i> |
| Pistachio | <i>Pistacia vera</i> |
| Walnut | <i>Juglans</i> spp. |

CONIFER TREES

PS Clethodim can be used to control labeled grasses in Christmas tree farms, conifer nurseries, and conifer plantations (but not in forests).

| COMMON NAME | SCIENTIFIC NAME |
|---------------------------|------------------------------|
| Arborvitae, American | <i>Thuja occidentalis</i> |
| Cedars | <i>Cedrus</i> spp. |
| Cypress | <i>Taxodium</i> spp. |
| Fir, Douglas | <i>Pseudotsuga menziesii</i> |
| Firs | <i>Abies</i> spp. |
| Hemlock, Canadian/Eastern | <i>Tsuga Canadensis</i> |
| Hemlock, Western | <i>Tsuga heterophylla</i> |
| Pines | <i>Pinus</i> spp. |
| Spruces | <i>Picea</i> spp. |
| Yew | <i>Taxus</i> spp. |

NON-CROP OR NON-PLANTED AREAS

The following areas are considered non-crop or non-planted areas: rights-of-way including railroads, highways, roads, dividers, medians, pipelines, public utility lines, pumping stations, transformer stations and substations. Around airports, electric utilities, commercial buildings, manufacturing plants, storage yards, rail yards, fence lines, parkways, and post-harvest croplands. Also beneath greenhouse benches and around golf courses.

ANNUAL GRASSES

(EXCEPT FOR IN ESTABLISHED ALFALFA AND MINT)

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- Do not apply more than 8 fl. oz./A of PS Clethodim **per application** to the following crops: asparagus, carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beets, green onion, head and stem brassica vegetables, herbs, hops, leafy brassica greens, leafy greens, non-bearing food crops, root vegetables, safflower, sesame and strawberry. Do not apply more than 6 fl. oz./A of PS Clethodim per application to canola or mustard seed.

Annual Grasses (cont'd.)

| GRASS SPECIES | SCIENTIFIC NAME | WEED HEIGHT* (inches) | RATE FL. OZ./ACRE | HIGH RATE⁽¹⁾ |
|------------------------|------------------------------------|----------------------------------|--------------------------|--------------------------------|
| Barnyardgrass | <i>Echinochloa crus-galli</i> | 2 to 8 | 6 | 8 |
| Broadleaf Signalgrass | <i>Brachiaria platyphlla</i> | 2 to 6 | 6 | 8 |
| Brome | | | | |
| California | <i>Bromus carinatus</i> | 2 to 6 | 6 | 8 |
| Cheat | <i>Bromus secalinus</i> | 2 to 6 | 6 | 8 |
| Downy | <i>Bromus tectorum</i> | 2 to 6 | 6 | 8 |
| Ripgut | <i>Bromus diandrus</i> | 2 to 6 | 6 | 8 |
| Canarygrass | <i>Phalaris canariensis</i> | 1 to 4 | 6 | 8 |
| Crabgrass | | | | |
| Hairy | <i>Digitaria adscendens</i> | 2 to 6** | 6 | 8 |
| Large | <i>Digitaria sanguinalis</i> | 2 to 6** | 6 | 8 |
| Smooth | <i>Digitaria ischaemum</i> | 2 to 6** | 6 | 8 |
| Southern | <i>Digitaria ciliaris</i> | 2 to 6** | 6 | 8 |
| Crowfootgrass | <i>Dactyloctenium aegyptium</i> | 2 to 6** | 6 | 8 |
| Fall Panicum | <i>Panicum dichotomiflorum</i> | 2 to 8 | 6 | 8 |
| Field Sandbur | <i>Cenchrus incertus</i> | 2 to 6 | 6 | 8 |
| Foxtail | | | | |
| Giant | <i>Setaria feberi</i> | 2 to 12 | 6 | 8 |
| Green | <i>Setaria viridis</i> | 2 to 8 | 6 | 8 |
| Yellow | <i>Setaria glauca</i> | 2 to 8 | 6 | 8 |
| Goosegrass | <i>Eleusine indica</i> | 2 to 6** | 6 | 8 |
| Itchgrass | <i>Rottboellia cochinchinensis</i> | 2 to 6 | 6 | 8 |
| Junglerice | <i>Echinochloa colona</i> | 2 to 6 | 6 | 8 |
| Lovegrass (Stinkgrass) | <i>Eragrostis cilianensis</i> | 2 to 6 | 6 | 8 |
| Rabbitsfootgrass | <i>Polyopogon monspeliensis</i> | 1 to 4 | 6 | 8 |
| Red Rice | <i>Oryza sativa</i> | 1 to 3 | 6 | 8 |
| Ryegrass | | | | |
| Hardy | <i>Lolium remotum</i> | 2 to 6 | 6 | 8 |

Annual Grasses (cont'd.)

| GRASS SPECIES | SCIENTIFIC NAME | WEED HEIGHT* (inches) | RATE FL. OZ./ACRE | HIGH RATE⁽¹⁾ |
|--------------------------------------|--------------------------------|----------------------------------|--------------------------|--------------------------------|
| Italian | <i>Lolium multiflorum</i> | 2 to 6 | 6 | 8 |
| Seedling Johnsongrass | <i>Sorghum halepense</i> | 4 to 10 | 6 | 8 |
| Shattercane | <i>Sorghum bicolor</i> | 6 to 18 | 6 | 8 |
| Southwestern Cupgrass | <i>Eriochloa gracilis</i> | 2 to 6 | 6 | 8 |
| Sprangle top | | | | |
| Amazon | <i>Leptochloa panicoides</i> | 2 to 6 | 6 | 8 |
| Bearded | <i>Leptochloa fascicularis</i> | 2 to 6 | 6 | 8 |
| Mexican | <i>Leptochloa uninervia</i> | 2 to 6 | 6 | 8 |
| Red | <i>Leptochloa filiformis</i> | 2 to 6 | 6 | 8 |
| Texas Panicum | <i>Panicum texanum</i> | 2 to 6 | 6 | 8 |
| Volunteer Cereals ⁽²⁾ | | | | |
| Barley | <i>Hordeum vulgare</i> | 2 to 6 | 6 | 8 |
| Oats | <i>Avena sativa</i> | 2 to 6 | 6 | 8 |
| Rye | <i>Secale cereal</i> | 2 to 6 | 6 | 8 |
| Wheat | <i>Triticum aestivum</i> | 2 to 6 | 6 | 8 |
| Volunteer Corn ⁽³⁾ | <i>Zea mays</i> | 4 to 12 | 4 | 6 |
| Volunteer Corn ⁽³⁾ | <i>Zea mays</i> | 12 to 24 | 6 | 8 |
| Volunteer Corn (S.R.) ⁽⁴⁾ | <i>Zea mays</i> | 4 to 12 | 8 (suppression only) | |
| Volunteer Grain Sorghum | <i>Sorghum bicolor</i> | 8 to 12 | 6 | 8 |
| Wild Oats | <i>Avena fatua</i> | 2 to 6 | 6 | 8 |
| Wild Proso Millet | <i>Panicum miliaceum</i> | 2 to 10 | 6 | 8 |
| Witchgrass | <i>Panicum capillare</i> | 2 to 8 | 6 | 8 |
| Woolly Cupgrass | <i>Eriochloa villosa</i> | 2 to 8 | 6 | 8 |

* Generally occurs between 3-leaf stage and tillering.

** Length of lateral growth.

⁽¹⁾ Rates higher than 8 fl. oz./A may be applied in certain geographic areas, cropping situations, or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 8 to 16 fl. oz./A may be applied. Do not apply more than 8 fl. oz./A of PS Clethodim per application to the following crops: asparagus, carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beet, green onion, head and stem brassica vegetables, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, non-bearing food crops, root

vegetables, safflower, sesame and strawberry.

- (2) When a cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum PS Clethodim use rate for control is 8 fl. oz./A.
- (3) Includes Roundup Ready®, Liberty Link®, and IMI-CORN® volunteer corn.
- (4) Sethoxydim resistant volunteer corn.

| ANNUAL & PERENNIAL GRASS CONTROL IN ESTABLISHED ALFALFA AND MINT WITH PS CLETHODIM | | | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|----------------------|-----------|
| GRASS SPECIES | WEED STAGE | RATE FL. OZ./ACRE | HIGH RATE |
| Annual & Perennial Grasses Listed in Grass Table | See Table | 10 | 16 |
| <p>Mowing: The best control of annual grasses can be achieved by applying PS Clethodim before grass weeds are mowed. Once grass is mowed it becomes tougher to control, as much of the available leaf surface has been removed. In areas without a killing frost, some annuals can over-winter after having been mowed multiple times. These grasses form large crowns and may contain many viable buds. These grasses, even though they may be an annual grass, may require repeated applications of PS Clethodim for partial or complete control.</p> <p>Irrigated Alfalfa and Mint: Irrigation practices can be very critical to the successful use of PS Clethodim in established alfalfa and mint and may be necessary to initiate active growth of the weeds prior to application. Generally applications 2 to 4 days after irrigation are most effective. Irrigation made shortly after application (2 days) can be effective, but more consistent grass control occurs when the irrigation is made before the application.</p> <p>Aerial Application: Apply PS Clethodim in a minimum of 10 GPA in established alfalfa and mint when applying by air.</p> <p>Annual Grass Control: Apply PS Clethodim at the grass sizes indicated in the Annual Grass Table and rates indicated. If a grass has been cut, apply PS Clethodim after active growth has resumed and regrowth has reached the minimum height and before it reaches the maximum height indicated. Apply before the alfalfa/mint canopy covers the grasses and interferes with the spray coverage. Some annual grasses are spring- and summer-germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to PS Clethodim may vary from region to region. Also some annuals germinate over an extended period of time, and because control of small grasses is desired, applications after each weed flush may be required. As a general rule spray spring- and summer-germinating grasses as early in the season as possible, after initial green-up. Spray fall-germinating weeds in the fall soon after they begin growing but before any damage is done due to frost. Late fall applications may be less effective due to environmental conditions, such as frost, slower plant growth, or the onset of flowering.</p> <p>Perennial Grass Control: PS Clethodim effectively controls perennial grasses such as Bermudagrass, Johnsongrass, quackgrass, wirestem muhly, tall fescue, foxtail barley and orchardgrass. Due in part to lack of tillage, perennial grasses are more difficult to control in a perennial crop such as established alfalfa or mint. A program of repeated applications is usually necessary for best results. The best way to control perennial grasses is to do so in the year of stand establishment before rhizomes and stolons become large and difficult to kill.</p> <p>Use the high rate under heavy grass pressure and/or when grasses are at or near maximum height.</p> <p>Always add a crop oil concentrate at 1 qt./A by ground or 1% v/v (but not less than 1 pt./A) to the finished spray volume by air.</p> | | | |

ANNUAL BLUEGRASS CONTROL WITH PS CLETHODIM

| GRASS SPECIES | WEED STAGE | RATE FL. OZ./ACRE | HIGH RATE |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|------------------------------|------------------|
| Annual Bluegrass (<i>Poa annua</i>) | to 4-leaf | 6* | 16 |
| <ul style="list-style-type: none"> • Apply under favorable soil moisture and humidity, which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s). • Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass. • Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature. • Always add a crop oil concentrate at 1 qt./A by ground to the finished spray volume. *Use a minimum of 10 fl. oz./A to control annual bluegrass in seedling and established alfalfa and mint. | | | |

FOR USE IN ROUNDUP READY FIELD CORN (BURNDOWN)

| GRASS SPECIES | WEED SIZE (inches) | Rate when applied alone or with glyphosate |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------|-------------------------------------------------------|
| Field Corn | Up to 12 | 3 fl. oz./A |
| <ul style="list-style-type: none"> • For control of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting corn. • Care must be taken to avoid in-field boom (spray) overlaps or excessive crop injury may occur. • Replant no sooner than 6 days after application. • Adjuvant recommendations: NIS at 0.25% v/v plus AMS at 2.5 to 4 lbs./A. • Do not use COC or MSO with PS Clethodim in this use pattern. | | |

DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES IN CANOLA, DRY SHELLED BEAN & PEA (INCLUDING SOYBEAN), EDIBLE PODED LEGUME VEGETABLES, FLAX, MUSTARD SEED, SUCCULENT BEAN & PEA AND SUGAR BEET (REDUCED RATE DIRECTIONS NOT FOR USE IN CALIFORNIA)

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low or high temperatures and/or under very low humidity.

| GRASS SPECIES | SCIENTIFIC NAME | WEED HEIGHT (inches) | RATE FL. OZ./ACRE ⁽¹⁾ |
|-----------------------|--------------------------------|----------------------|-------------------------------------|
| Barnyardgrass | <i>Echinochloa crus-galli</i> | 1 to 4 | 4 |
| Broadleaf Signalgrass | <i>Brachiaria platyphlla</i> | 1 to 4 | 5 |
| Crabgrass | | | |
| Large | <i>Digitaria sanguinalis</i> | 1 to 3* | 4 |
| Large | <i>Digitaria sanguinalis</i> | 1 to 4* | 5 |
| Smooth | <i>Digitaria ischaemum</i> | 1 to 3* | 4 |
| Smooth | <i>Digitaria ischaemum</i> | 1 to 4* | 5 |
| Southern | <i>Digitaria ciliaris</i> | 1 to 4* | 5 |
| Fall Panicum | <i>Panicum dichotomiflorum</i> | 1 to 4 | 4 |
| Foxtail | | | |
| Giant | <i>Setaria faberi</i> | 1 to 4 | 4 |
| Green | <i>Setaria viridis</i> | 1 to 4 | 4 |
| Millet | <i>Setaria italica</i> | 1 to 4 | 5 |
| Yellow | <i>Setaria glauca</i> | 1 to 4 | 4 |
| Seedling Johnsongrass | <i>Sorghum halepense</i> | 1 to 6 | 5 |
| Shattercane | <i>Sorghum bicolor</i> | 4 to 10 | 4 |
| Texas Panicum | <i>Panicum texanum</i> | 1 to 4 | 5 |

| GRASS SPECIES | SCIENTIFIC NAME | WEED HEIGHT (inches) | RATE FL. OZ./ACRE ⁽¹⁾ |
|-------------------|--------------------------|----------------------|-------------------------------------|
| Volunteer Cereals | | | |
| Barley | <i>Hordeum vulgare</i> | 1 to 4 | 5 |
| Oats | <i>Avena sativa</i> | 1 to 4 | 5 |
| Wheat | <i>Triticum aestivum</i> | 1 to 4 | 5 |
| Volunteer Corn** | <i>Zea mays</i> | 4 to 12 | 4 |
| Wild Proso Millet | <i>Panicum miliaceum</i> | 1 to 6 | 4 |
| Wild Oats | <i>Avena fatua</i> | 1 to 4 | 5 |

* Length of lateral growth

** Not S.R. Corn

⁽¹⁾ Always add a crop oil concentrate at 1 qt./A by ground application to the finished spray volume.

PERENNIAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- Do not apply more than 8 fl. oz./A of PS Clethodim per application to the following crops: asparagus, carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beet, green onion, head & stem brassica vegetables, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, non-bearing food crops, root vegetables, safflower, sesame and strawberry.
- Do not apply more than 6 fl. oz./A of PS Clethodim per application to canola or mustard seed.

| GRASS SPECIES | WEED HEIGHT (inches) | RATE FL. OZ./ACRE | HIGH RATE |
|--------------------------------------------|-------------------------|----------------------|-----------|
| <i>Bermudagrass (Cynodon dactylon)</i> | | | |
| First Application | 3 (or up to 6" runners) | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 3 (or up to 6" runners) | 8 | 16 |
| <i>Fescue, Tall (Festuca arundinacea)</i> | | | |
| First Application | 4 to 8 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 4 to 8 | 8 | 16 |
| <i>Foxtail Barley (Hordeum jubatum)</i> | | | |
| First Application | 2 to 6 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 2 to 6 | 8 | 16 |
| <i>Orchardgrass (Dactylis glomerata)</i> | | | |
| First Application | 4 to 8 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 4 to 8 | 8 | 16 |

Perennial Grasses (cont'd.)

| GRASS SPECIES | WEED HEIGHT (inches) | RATE FL. OZ./ACRE | HIGH RATE |
|---------------------------------------------------|-----------------------------|------------------------------|------------------|
| Quackgrass* (<i>Elytrigia repens</i>) | | | |
| First Application | 4 to 12 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 4 to 12 | 8 | 16 |
| Rhizome Johnsongrass (<i>Sorghum halepense</i>) | | | |
| First Application | 12 to 24 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 6 to 18 | 6 | 8 |
| Wirestem Muhly (<i>Muhlenbergia frondosa</i>) | | | |
| First Application | 4 to 8 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 4 to 8 | 8 | 16 |
| Perennial Bluegrass* | | | |
| Roughstalk (<i>Poa trivialis</i>) | | | |
| Kentucky (<i>Poa pratensis</i>) | | | |
| First Application | 2 to 4 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 2 to 4 | 8 | 16 |
| Bentgrass* (<i>Agrostis</i> spp.) | | | |
| First Application | 2 to 4 | - | 16 |
| Repeat Application(s) (if regrowth occurs) | 2 to 4 | - | 16 |

*Control of quackgrass, perennial bluegrass and bentgrass with PS Clethodim may be enhanced by adding AMS at 2.5 to 4.0 lbs./A.

TANK MIXES

INFORMATION

The labels for each of the herbicides recommended for tank mixing with PS Clethodim are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the PS Clethodim label in certain considerations. Those concerns may include, but are not limited to:

1. Geographic restrictions – all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (e.g. protective clothing, reentry time, posting);
5. Soil type or soil characteristics (e.g. pH, OM);
6. Maximum dosage or number of applications per season;
7. Rain free period required; or
8. Application timing (e.g. pre-harvest interval)
9. Do not exceed the total season rates.

THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.

TANK MIX APPLICATION OF PS CLETHODIM AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label.
- Apply when the first grass or broadleaf weed species in a mixed population reaches the recommended height or growth stage for treatment.
- Apply under favorable soil moisture and humidity that exist a few days after rainfall or within seven days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate recommended for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of PS Clethodim, as specified in the respective size and rate tables.
- Do not tank mix PS Clethodim when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of PS Clethodim. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing PS Clethodim with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. **Agitation should continue until all spray solution has been applied.**

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control. Mixing and compatibility qualities should be verified by a jar test.

INFORMATION ON ANTAGONISM

Tank mixes of PS Clethodim with postemergence broadleaf herbicides have shown some reduction or failure to control certain

grass species which would have otherwise been controlled when PS Clethodim is applied alone. Activity of the postemergence broadleaf herbicide in the tank mix is not affected.

ALFALFA

Table 1. PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR ALFALFA
(Refer to tables above for specific grasses and growth stages)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|---------------------------------------------------------------------------------------|-------------------------------------------------------------------|------------------------------------------------|-------------------------------------------|------|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| | | | | |
| PS Clethodim + 2,4-DB ⁽⁴⁾ | 10 to 16 fl. oz. + Refer to 2,4-DB label | 10 to 16 fl. oz. + Refer to 2,4-DB label | 1% | 1% |
| PS Clethodim + PURSUIT DG ⁽⁵⁾ or PURSUIT ⁽⁵⁾ | 10 to 16 fl. oz. + 1.08 to 2.16 oz. or 3 to 6 fl. oz. | - | 1% | 1% |
| PS Clethodim + BUCTRIL® 2L ⁽⁶⁾ or BUCTRIL GEL ^(6,7) | 10 to 16 fl. oz. + 1.0 to 1.5 pts. or 0.5 to 0.75 pt. | - | 0.5% | 0.5% |

- ⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- ⁽²⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- ⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- ⁽⁴⁾ PS Clethodim plus 2,4-DB may increase the severity of crop injury when tank mixed. Alfalfa plants will generally outgrow this temporary crop injury within a few weeks.
- ⁽⁵⁾ Before using this tank mix, read and understand the PURSUIT or PURSUIT DG labels for geographical restrictions and restrictions regarding alfalfa growth stage and type. Failure to do so can result in crop injury to alfalfa. Do not feed, graze, or harvest alfalfa for 30 days following an application of PURSUIT to alfalfa.
- ⁽⁶⁾ In the states of Washington, Oregon, Idaho, Montana, Wyoming, Colorado, Utah, Nevada and the western halves of North Dakota, South Dakota, Nebraska, and Kansas: The PS Clethodim plus BUCTRIL or BUCTRIL GEL tank mix must be applied in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 2 trifoliates. Unacceptable crop injury may occur to alfalfa seedlings less than the 2 trifoliolate leaf stage. PS Clethodim plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 80°F at (and 3 days following) application can result in unacceptable crop injury. In the states not listed above, apply in the fall or spring to seedling alfalfa when the majority of the field has a minimum

of 4 trifoliolate leaves. When the alfalfa stand is uneven and conditions favor leaf burn, unacceptable crop injury may occur to alfalfa in the 2 trifoliolate or smaller stage of growth. PS Clethodim plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 70°F at (and 3 days following) application can result in unacceptable crop injury. Crop leaf burn can occur following PS Clethodim plus BUCTRIL or BUCTRIL GEL application. Warm, humid conditions may enhance leaf burn. New crop growth will not be affected.

- ⁽⁷⁾ Do not apply when alfalfa is under moisture, temperature, insect or disease stress or has been stressed by other pesticide carryover or application.

CANOLA

Table 2. REDUCED RATE PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR CANOLA

(Refer to the tables above for specific grasses and growth stages.)

| PRODUCT | APPLICATION RATES/ACRE | | AMMONIUM SULFATE | |
|------------------------------------------------------------|-----------------------------------|-------------------|------------------|----------|
| | ANNUAL GRASSES ⁽¹⁾ | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim ⁽²⁾ + LIBERTY ⁽³⁾ | 4 to 5 fl. oz. + 34 fl. oz. | - | 3.0 lbs. | 3.0 lbs. |

⁽¹⁾ Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table.

⁽²⁾ Do not apply PS Clethodim tank mix during or after bolting or flowering or crop injury may occur.

⁽³⁾ For use only on LibertyLink® canola.

COTTON

Table 3. PS CLETHODIM TANK MIXED WITH COBRA® AND MSMA APPLIED POST DIRECTED TO COTTON

| PRODUCT ⁽¹⁾ | APPLICATION RATES/ACRE ⁽²⁾ | | CROP OIL CONCENTRATE ⁽³⁾ V/V | COMMENTS |
|------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|--------------------------------------------|------------------------------------------------------------------------|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | |
| PS Clethodim ⁽⁴⁾ + COBRA + MSMA (4.0 lbs./gal.) or MSMA (6.6 lbs./gal.) | 6 to 8 fl. oz. | 8 to 16 fl. oz. | 1% | Reduce broadcast rate in proportion to the band area actually treated. |
| | See COBRA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the PS Clethodim label for weed height and species controlled. | | | |
| | See MSMA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the PS Clethodim label for weed height and species controlled. | | | |

- (1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- (4) If at the time of application, grass height is so tall that post-directed applications cannot get good coverage over the top of the grassy weeds, then poor control may result and a second (non-post directed) application of PS Clethodim may be necessary.

Table 4. PS CLETHODIM TANK MIXED WITH BUCTRIL 4 EC TO CONTROL EMERGED WEEDS IN BXN COTTON AS A BROADCAST APPLICATION

| PRODUCT ⁽¹⁾ | APPLICATION RATES/ACRE ⁽²⁾ | CROP OIL CONCENTRATE PER ACRE ⁽³⁾ | COMMENTS ⁽⁷⁾ |
|------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------|----------------------------------------------|------------------------------------|
| | ANNUAL GRASSES | | |
| PS Clethodim + BUCTRIL 4 EC ^(4,5,6) | 8 to 16 fl. oz. + See BUCTRIL 4 EC label for rates to control broadleaf weeds and height limitations for cotton | 1 qt. | See charts for grasses controlled. |

- (1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage.
- (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim at the recommended rate with the appropriate amount of crop oil concentrate in a non-BUCTRIL tank mix.
- (3) Always add a crop oil concentrate at 1 qt./A by ground in the finished spray solution.
- (4) Applications of BUCTRIL 4 EC can be made only to cotton that has been genetically modified for crop tolerance to postemergence over-the-top application of bromoxynil.
- (5) Do not apply the PS Clethodim plus BUCTRIL tank mix within 75 days of harvest.
- (6) Do not exceed 2 applications of BUCTRIL before cotton is 12 inches tall and one application after 12 inches tall.
- (7) Use a minimum of 10 gals. of spray solution per acre.

Table 5. PS CLETHODIM TANK MIXED WITH GLYPHOSATE TO CONTROL EMERGED GRASSES IN COTTON AS A BROADCAST APPLICATION

| PRODUCT | APPLICATION RATES/ACRE ⁽¹⁾ | | ADJUVANT | | COMMENTS |
|---------------------------------|----------------------------------------------------------------------------------------------|-------------------|-----------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| | ANNUAL GRASSES | PERENNIAL GRASSES | Glyphosate formulation with built in adjuvant | Glyphosate formulation without built in adjuvant | |
| PS Clethodim + GLYPHOSATE | 6 to 8 fl. oz. | 8 to 16 fl. oz. | Non-ionic surfactant @ 0.125 to 0.25% v/v plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier | Crop oil concentrate @ 1 pt./A plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier | See charts for grasses controlled. Use a minimum of 10 gals of spray solution per acre. |
| | See glyphosate label for rates to control broadleaf weeds and height limitations for cotton. | | | | |

⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim at the recommended rate with the appropriate amount of crop oil concentrate.

DRY SHELLED AND SUCCULENT BEANS

Table 6. PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR DRY SHELLED AND SUCCULENT BEANS (Refer to the tables above for specific grasses and growth stages.)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|--------------------------------|-----------------------------------------|--------------------------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + BASAGRAN® | 8 to 10 fl. oz. + 1.0 to 2.0 pts. | 10 to 16 fl. oz. + 1 to 2 pts. | 1% | 1% |

⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.

⁽²⁾ Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

FLAX

**Table 7. REDUCED RATE PS CLETHODIM TANK MIXED WITH BROADLEAF HERBICIDES FOR FLAX
(Refer to the tables above for specific grasses and growth stages)**

| PRODUCT | APPLICATION RATES/ACRE | | CROP OIL CONCENTRATE | |
|------------------------------------------------------------|----------------------------------------|-------------------|----------------------|-------|
| | ANNUAL GRASSES ⁽¹⁾ | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + BRONATE ADVANCED™ ^(2,3) | 4 to 5 fl. oz. + 11.4 fl. oz. | - | 1 pt. | 1 pt. |
| PS Clethodim + BRONATE® ^(2,3) | 4 to 5 fl. oz. + 0.9 pt. | - | 1 pt. | 1 pt. |
| PS Clethodim + BUCTRIL ^(2,3) | 4 to 5 fl. oz. + 1.0 pt. | - | 1 pt. | 1 pt. |
| PS Clethodim + RHONOX® ^(2,3) | 4 to 5 fl. oz. + 0.25 to 0.5 pt. | - | 1 pt. | 1 pt. |

⁽¹⁾ Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table.

⁽²⁾ Do not apply PS Clethodim tank mix during or after the bud stage or to ornamental flax or crop injury may occur.

⁽³⁾ Do not apply tank mixes if temperatures are expected to exceed 85°F at (or 3 days following) application or crop injury may occur.

SOYBEAN

Table 8. PS CLETHODIM TANK MIXES⁽³⁾ TO CONTROL ANNUAL GRASSES WHEN USED AS A BURNDOWN IN NO-TILL SOYBEAN

| PRODUCT | PRODUCT RATE/ACRE ⁽¹⁾ | GRASS HEIGHT (inches) | CROP OIL CONCENTRATE /ACRE ⁽²⁾ | 28%N OR 32%N QTS./A OR 2.5 TO 4.0 LBS. AMS |
|-------------------------------------------|-----------------------------------|----------------------------------------|-------------------------------------------|--------------------------------------------|
| PS Clethodim + 2,4-D ester ⁽³⁾ | 3 fl. oz. | Foxtail 1 to 3 Fall Panicum 1 to 3 | 1 qt. | 1 to 2 qts. or 2.5 to 4.0 lbs. AMS |
| | 4 fl. oz. | Foxtail 1 to 4 Fall Panicum 1 to 4 | 1 qt. | 1 to 2 qts. or 2.5 to 4.0 lbs. AMS |
| | 6 to 8 fl. oz. + 0.5 lb. ai | (See Grass Chart for grasses claimed.) | 1 qt. | 1 to 2 qts. or 2.5 to 4.0 lbs. AMS |

* 2,4-D ester should not be used where drift sensitive crops may be grown.

- ⁽¹⁾ If regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim according to the appropriate size and rate recommendations.
- ⁽²⁾ Always use a crop oil concentrate at the listed rate in the finished spray volume.
- ⁽³⁾ The following products can be tank mixed with PS Clethodim plus 2,4-D ester: VALOR®, AUTHORITY® BROADLEAF, CANOPY XL®, DUAL® 8 E, DUAL II®, DUAL MAGNUM®, PROWL®, SENCOR®, SENCOR plus the DUAL products and TURBO®.

Table 9. PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN
(Refer to the tables above for specific grasses and growth stages.)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|------------------------------|---------------------------------------|--------------------------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + COBRA | 6 to 8 fl. oz. + 12.5 fl. oz. | 8 to 16 fl. oz. + 12.5 fl. oz. | 0.5 to 1% | 1% |
| PS Clethodim + BASAGRAN 4 SL | 8 to 10 fl. oz. + 1 to 2 pts. | 10 to 16 fl. oz. + 1 to 2 pts. | 1% | 1% |

Soybean (cont'd.)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|-------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|-------------------------------------------|-------------------|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + Glyphosate (For use on Roundup Ready soybean only) | 6 to 8 fl. oz. + 0.75 to 3.0 lb. ai | 8 to 16 fl. oz. + 0.75 to 3.0 lb. ai | 0.5 to 1% ⁽⁴⁾ | 1% ⁽⁴⁾ |
| PS Clethodim + ULTRA BLAZER [®] | 6 to 8 fl. oz. + 1 to 1.5 pts. | 6 to 8 fl. oz. + 1 to 1.5 pts. | 0.5% to 1% | 1% |
| PS Clethodim + FLEXSTAR [®] HL ⁽⁶⁾ | 6 to 8 fl. oz. + Refer to the FLEXSTAR HL label for specific application rates. | 8 to 16 fl. oz. + Refer to the FLEXSTAR HL label for specific application rates. | 1% | 1% |
| PS Clethodim + CLASSIC [®] 25 DG | 8 to 10 fl. oz. + 0.5 to 0.75 oz. | 10 to 16 fl. oz. + 1.44 oz. | 1% | 1% |
| PS Clethodim ⁽⁴⁾ + PURSUIT [®] 70 DG | 6 to 8 fl. oz. + 1.44 oz. | 8 to 16 fl. oz. + 1.44 oz. | 1% | 1% |
| PS Clethodim ⁽⁵⁾ + COBRA + CLASSIC 25 DG | 8 to 10 fl. oz. + 6 to 8 fl. oz. + 0.5 to 0.75 oz. | - | 0.5% | 1% |

Soybean (cont'd.)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|-----------------------------------------------------------------------------------|---------------------------------------------------------------|-------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim ⁽⁵⁾ + COBRA + BASAGRAN 4 SL | 8 to 10 fl. oz. + 6 to 10 fl. oz. + 1 to 1.5 pts. | - | 0.5% | 1% |
| PS Clethodim ⁽⁵⁾ + COBRA + PURSUIT 70 DG | 8 to 10 fl. oz. + 6 to 10 fl. oz. + 1.44 oz. | - | 0.5% | 1% |
| PS Clethodim ⁽⁵⁾ + STORM ⁽⁶⁾ | 8 to 10 fl. oz. + 1.5 pts. | - | 0.5% | 1% |
| PS Clethodim ⁽⁵⁾ + RESOURCE ⁽⁶⁾ + PURSUIT 70 DG | 8 to 10 fl. oz. + 4 fl. oz. + 1.44 oz. | - | 1% | 1% |
| PS Clethodim ⁽⁵⁾ + RESOURCE + BASAGRAN | 8 to 10 fl. oz. + 4 fl. oz. + 1 pt. | - | 1% | 1% |
| PS Clethodim ⁽⁵⁾ + RESOURCE + CLASSIC | 8 to 10 fl. oz. + 4 fl. oz. + 0.5 oz. | - | 1% | 1% |

Soybean (cont'd.)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|----------------------------------------------------------------------------------------|----------------------------------------------------------------------|--------------------------------------------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim ⁽⁵⁾ + COBRA + RESOURCE | 6 to 8 fl. oz. + 6 fl. oz. + 4 fl. oz. | - | 0.5% | 1% |
| PS Clethodim ⁽⁵⁾ + FIRSTRATE [®] | 6 to 8 fl. oz. + 0.3 oz. | 8 to 16 fl. oz. + 0.3 oz. | 1% | - |
| PS Clethodim ⁽⁵⁾ + COBRA + FIRSTRATE | 6 to 8 fl. oz. + 6 to 8 fl. oz. + 0.3 oz. | 8 to 16 fl. oz. + 6 to 8 fl. oz. + 0.3 oz. | 1% | - |
| PS Clethodim ⁽⁵⁾ + RAPTOR [®] (1 AS) | 6 to 8 fl. oz. + 4 to 5 fl. oz. | - | 1% | - |
| PS Clethodim ⁽⁵⁾ + COBRA + RAPTOR (1 AS) | 6 to 8 fl. oz. + 6 to 8 fl. oz. + 4 to 5 fl. oz. | - | 1% | - |
| PS Clethodim ⁽⁵⁾ + SYNCHRONY [®] STS [™] | 6 to 8 fl. oz. ⁽⁷⁾ + 0.5 oz./A | - | 1 qt. | - |
| PS Clethodim ⁽⁵⁾ + COBRA Herbicide + SYNCHRONY STS [™] | 6 to 8 fl. oz. ⁽⁷⁾ + 4 to 8 fl. oz. + 0.5 oz. | - | 1 pt. | - |

Soybean (cont'd.)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|-------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim ⁽⁵⁾ + RESOURCE | 6 to 8 fl. oz. + 4 to 12 fl. oz. | - | 1 qt. | - |
| PS Clethodim ⁽⁵⁾ + FRONTROW™ | 8 to 10 fl. oz. + Refer to FRONTROW label for use rates. | - | 1% | - |
| PS Clethodim + FIRSTRATE + FLEXSTAR HL ⁽⁶⁾ | 6 to 8 fl. oz. + 0.3 oz. + Refer to the FLEXSTAR HL label for specific application rates. | 8 to 16 fl. oz. + 0.3 oz. + Refer to the FLEXSTAR HL label for specific application rates. | 1% | - |

- ⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- ⁽²⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- ⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- ⁽⁴⁾ The addition of 2.5 lb. of ammonium sulfate is required when PS Clethodim is tank mixed with glyphosate. If the glyphosate formulation has a stand alone built in adjuvant, add 0.125% v/v non-ionic surfactant in place of crop oil concentrate. If the glyphosate formulation does not have a built in adjuvant, add 0.5 to 1% crop oil concentrate for ground application and 1% v/v for aerial application.
- ⁽⁵⁾ The addition of 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N) is recommended when PS Clethodim is tank mixed with PURSUIT, RESOURCE, STORM, FIRSTRATE, SYNCHRONY, RAPTOR, FRONTROW, COBRA plus CLASSIC, COBRA plus BASAGRAN, COBRA plus PURSUIT, COBRA plus FIRSTRATE, COBRA plus SYNCHRONY, and COBRA plus RAPTOR. An equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.
- ⁽⁶⁾ Refer to FLEXSTAR HL label for geographic and rotational restrictions.
- ⁽⁷⁾ Annual grasses and sizes controlled with these tank mixes are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table.

Table 10. REDUCED RATE PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN (Refer to table for reduced rate use in canola, dry shelled bean & pea, edible podded legume vegetables, flax, mustard seed, soybean, succulent bean & pea and sugar beet for small annual grasses for specific grasses and growth stages)

| PRODUCT | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ^(3,4) (V/V) | |
|------------------------------------|---------------------------------------|-------------------|---------------------------------------------|-----|
| | ANNUAL GRASSES ⁽²⁾ | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + FIRSTRATE | 4 to 8 fl. oz. + 0.3 oz. | - | 1% | 1% |
| PS Clethodim + PURSUIT 70 DG | 4 to 6 fl. oz. + 1.44 oz. | - | 1% | 1% |

⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.

⁽²⁾ Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table.

⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

⁽⁴⁾ The addition of 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N) is required when PS Clethodim is tank mixed at reduced rates. An equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.

PEANUT (INCLUDING PERENNIAL)

Table 11. PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR PEANUT (INCLUDING PERENNIAL)
(Refer to the tables above for specific grasses and growth stages.)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|-----------------------------------|-----------------------------------------|-------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES ⁽²⁾ | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + BASAGRAN | 8 to 10 fl. oz. + 1.0 to 2.0 pts. | - | 1% | 1% |
| PS Clethodim + ULTRA BLAZER | 8 to 10 fl. oz. + 0.5 to 1.5 pts. | - | 1% | 1% |
| PS Clethodim + STORM | 8 to 10 fl. oz. + 1.5 pts. | - | 1% | 1% |

- ⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- ⁽²⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- ⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

| GRASS SUPPRESSION FOR HARVEST EFFICIENCY IN PEANUT (INCLUDING PERENNIAL) WITH PS CLETHODIM | | | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------|-------------------|-----------|
| GRASS SPECIES | WEED STAGE | RATE FL. OZ./ACRE | HIGH RATE |
| Annual and perennial grasses that exceed height claimed for control on height charts "ANNUAL GRASSES" & "PERENNIAL GRASSES" | Up to and including grasses in the seed head stage. | 16 | 32 |
| Do not apply as part of a tank mix when applying PS Clethodim for grass suppression. Add a crop oil concentrate at 1 qt./A by ground to the finished spray volume. | | | |

SUGAR BEET

Table 12. PS CLETHODIM TANK MIXED WITH STINGER® APPLIED TO SUGAR BEET
(Refer to the tables above for specific grasses and growth stages)

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) | |
|------------------------------|---------------------------------------|-------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + STINGER | 6 to 8 fl. oz. | 8 to 16 fl. oz. | 1% | 1% |
| | See STINGER label for rates. | | | |

- ⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- ⁽²⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- ⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

Table 13. PS CLETHODIM TANK MIXED WITH BETAMIX® or BETANEX® APPLIED TO SUGAR BEET

| PRODUCT ⁽¹⁾ | WEEDS CONTROLLED | | WEED HEIGHT (inches) | APPLICATION RATE/ ACRE ⁽²⁾ |
|--------------------------------------------------------------|-------------------|------------------------------|--------------------------------------------------------------------------------------------------------------|------------------------------------------|
| | COMMON NAME | SCIENTIFIC NAME | | |
| PS Clethodim ⁽³⁾ + BETAMIX or BETANEX | Barnyardgrass | <i>Echinochloa crusgalli</i> | 1 to 3 | 8 fl. oz. |
| | Foxtail | <i>Setaria</i> spp. | 1 to 3 | |
| | Foxtail Millet | <i>Setaria italic</i> | 1 to 3 | |
| | Wild Oat | <i>Avena fatua</i> | 1 to 3 | |
| | Wild Proso Millet | <i>Panicum miliaceum</i> | 1 to 3 | |
| | | | See BETAMIX label for rates to control broadleaf weeds. No additives are recommended in the tank mix. | |
| | | | See BETANEX label for rates to control broadleaf weeds. No additives are recommended in the tank mix. | |

⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

⁽²⁾ Do not use crop oil concentrate. No additives are recommended in the tank mix. If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.

⁽³⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim at full label rate with appropriate rate of crop oil concentrate.

Table 14. PS CLETHODIM PLUS BETANEX OR BETAMIX TANK MIX FOR THREE SEQUENTIAL APPLICATIONS FOR ANNUAL GRASS CONTROL (MICRO-RATE APPLICATION)

| PRODUCT | APPLICATION RATES/ACRE ⁽¹⁾ | | METHYLATED SEED OIL ⁽²⁾ (V/V) | |
|-----------------------------------------------|-----------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|------------------------------------------|------|
| | ANNUAL GRASSES | GRASSES CONTROLLED (inches) | GROUND | AIR |
| PS Clethodim + BETANEX or BETAMIX | 2 to 3 fl. oz. + 8 to 12 fl. oz. ⁽³⁾ or 8 to 12 fl. oz. ⁽³⁾ | Green Foxtail (1-2) Yellow Foxtail (1-2) Barnyardgrass (1-2) Wild Oat (1-2) Volunteer Cereals (1-2) | 1.5% | 1.5% |

⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

⁽²⁾ Always use a methylated seed oil at the listed rate (but not less than 1 pt./A) in the finished spray volume.

⁽³⁾ Use 8 fl. oz./A rate when sugar beet are in the cotyledon to 4 leaf stage. Rate can be increased up to 12 fl. oz./A when the smallest sugar beet plants in the field are in the 4 true leaf stage or larger.

Directions for Use for Micro-Rate Applications to Sugar Beet Application Information

Multiple micro-rate applications of PS Clethodim in tank mixtures with reduced rates of BETANEX or BETAMIX and methylated seed oils may be applied by air or ground equipment to sugar beet to control early germinating annual grasses listed above. The rate of BETANEX or BETAMIX must not exceed 0.12 lb. ai./A (broadcast application) when in combination with these spray adjuvants. Note that maximum rate allowed varies depending on crop growth stage. The use of wetting agents or spray adjuvants with conventional rates (0.73 to 1.22 lb. ai./A) or multiple low rate (0.24 to 0.73 lb. ai./A) applications of BETANEX or BETAMIX is prohibited on the BETANEX and BETAMIX master label. Favorable climatic conditions (good conditions for plant growth and development) are essential for adequate weed control. All use precautions and restrictions on the BETANEX and BETAMIX master labels must be followed.

Directions for Using Micro-Rate Multiple Applications of PS Clethodim Tank Mixes

Apply PS Clethodim in broadcast applications only at a rate of 2 to 3 fl. oz./A in tank mixture with either BETANEX or BETAMIX following the directions for use on the tank mix partner label. A minimum of three sequential applications of 2 fl. oz./A or a minimum of 2 sequential applications of 3 fl. oz./A should be utilized for PS Clethodim tank mixtures. A minimum of 3 sequential applications of BETAMIX or BETANEX should be used. Accurate timing is essential; make initial application immediately after weeds emerge, and make repeat applications on 5 to 7 day intervals. If weed control is not adequate due to climatic conditions, spray coverage or other factors, return to conventional application rates of PS Clethodim (6 to 8 fl. oz./A) and add rates of BETANEX or BETAMIX as directed on their label. When using conventional rates of BETANEX or BETAMIX in tank mixtures with PS Clethodim, a spray adjuvant is not recommended.

Use Precautions for Micro-Rate Applications: (See PS Clethodim, BETANEX and BETAMIX master label for further use precautions.)

Not all weeds will be adequately controlled, even with favorable climatic conditions. Conventional rates of PS Clethodim, BETANEX or BETAMIX and/or hand labor may be required if multiple micro-rate applications do not adequately control weeds. Plugging of spray nozzles may be encountered due to the potential for formation of a precipitate in the spray solution that is often associated with micro-rate applications. Prime Source, LLC will not be responsible for any nozzle plugging that may occur with the use of multiple micro-rate applications. Methylated seed oils must not be added if the BETANEX or BETAMIX rate exceeds 0.12 lb. ai/A broadcast, as the addition of methylated seed oils could increase the possibility of crop injury at dosage rates greater than 0.12 lb. ai/A.

GROUND APPLICATION

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 10 gals. and a maximum of 20 gals. of spray solution per acre. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

AERIAL APPLICATION

Use of sufficient spray volumes is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 15 gals. of spray solution per acre.

Table 15. TANK MIX APPLICATION OF PS CLETHODIM AND FUNGICIDES FOR CONTROL OF GRASS WEEDS AND DISEASES IN SUGAR BEET

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽³⁾ (V/V) |
|-------------------------------------------|---------------------------------------|------------------------------------|-------------------------------------------|
| | ANNUAL GRASSES | PERENNIAL GRASSES | |
| PS Clethodim + EMINENT [®] | 6 to 8 fl. oz. + 13 fl. oz. | 8 to 16 fl. oz. + 13 fl. oz. | 1% |

⁽¹⁾ If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix fungicide), according to the appropriate size and rate recommendations.

⁽²⁾ Refer to PS Clethodim and fungicide label for rates and weeds and diseases controlled.

⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

Table 16. TANK MIX APPLICATION OF PS CLETHODIM AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT, PEANUT (INCLUDING PERENNIAL), SOYBEAN AND SUNFLOWER

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE (V/V) ⁽³⁾ | CROP | | | | | |
|--------------------------------------------------------|------------------------------------------------------------|-------------------------------------------------------|-------------------------------------------|------------------------|--------|-----------------------|--------|---------|-----------|
| | ANNUAL GRASSES | PERENNIAL GRASSES | | Alfalfa ⁽⁶⁾ | Cotton | Mint ^(4,5) | Peanut | Soybean | Sunflower |
| PS Clethodim + ORTHENE [®] 75 S or ORTHENE 97 | 6 to 8 fl. oz. + 0.33 to 1.33 lbs. or 0.25 to 1.0 lb. | 6 to 8 fl. oz. + 0.33 to 1.33 lbs. or 0.25 to 1.0 lb. | 1% | | X | X | X | | |
| PS Clethodim + ORTHENE 90 S ⁽⁶⁾ | 6 to 8 fl. oz. + 0.25 to 1 lb. | 8 to 16 fl. oz. + 0.25 to 1 lb. | 1% | | X | X | X | X | |
| PS Clethodim + DANITOL [®] 2.4 EC | 6 to 8 fl. oz. + 10 2/3 to 16 fl. oz. | 8 to 16 fl. oz. + 10 2/3 to 16 fl. oz. | 1% | | X | | X | | |
| PS Clethodim + ASANA XL [®] | 6 to 8 fl. oz. + Refer to ASANA XL label | 8 to 16 fl. oz. + Refer to ASANA XL label | 1% | | | | | | X |
| PS Clethodim + WARRIOR [®] | 6 to 8 fl. oz. + Refer to WARRIOR label | 8 to 16 fl. oz. + Refer to WARRIOR label | 1% | | | | | | X |
| PS Clethodim + WARRIOR | 10 to 16 fl. oz. ⁽⁷⁾ + Refer to WARRIOR label | 10 to 16 fl. oz. + Refer to WARRIOR label | 1% | X | | | | | |
| PS Clethodim + BAYTHROID [®] | 10 to 16 fl. oz. ⁽⁷⁾ + Refer to BAYTHROID label | 10 to 16 fl. oz. + Refer to BAYTHROID label | 1% | X | | | | | |

| PRODUCT ⁽²⁾ | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE (V/V) ⁽³⁾ | CROP | | | | | |
|----------------------------|-------------------------------------------------------------------|-------------------------------------------------------------------|-------------------------------------------|------------------------|--------|-----------------------|--------|---------|-----------|
| | ANNUAL GRASSES | PERENNIAL GRASSES | | Alfalfa ⁽⁴⁾ | Cotton | Mint ^(4,9) | Peanut | Soybean | Sunflower |
| PS Clethodim + DIMETHOATE® | 10 to 16 fl. oz. ⁽⁷⁾ + Refer to DIMETHOATE label | 10 to 16 fl. oz. ⁽⁷⁾ + Refer to DIMETHOATE label | 1% | X | | | | | |
| PS Clethodim + LORSBAN® | 10 to 16 fl. oz. ⁽⁷⁾ + Refer to LORSBAN label | 10 to 16 fl. oz. + Refer to LORSBAN label | 1 to 2 pt. ⁽⁸⁾ | X | | | | | |
| PS Clethodim + POUNCE® | 10 to 16 fl. oz. ⁽⁷⁾ + Refer to POUNCE label | 10 to 16 fl. oz. + Refer to POUNCE label | 1% | X | | | | | |

⁽¹⁾ If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of PS Clethodim alone (without a tank mix fungicide), according to the appropriate size and rate recommendations.

⁽²⁾ Refer to PS Clethodim and fungicide label for rates and weeds and diseases controlled.

⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

⁽⁴⁾ Certain insecticides may cause temporary phytotoxic symptoms on alfalfa and mint foliage. Refer to the insecticide label for further information. It is suggested that prior to using any of these insecticides/herbicide tank mixtures, that a small area of the field be treated first and observations for crop injury be made prior to treating the whole field.

⁽⁵⁾ The PS Clethodim rate should be 6 to 8 fl. oz./A for annual grass control in baby mint, minimum of 8 fl. oz./A for annual grass control in established mint and 8 to 16 fl. oz./A for perennial grass control. Crop oil concentrate should be added at the rate of 1.0 to 2.0 pts./A.

⁽⁶⁾ Insecticide tank mix use with ORTHENE 90 S in soybean is permitted only in a state having an approved Section 24(c) registration for ORTHENE 90 S use in soybean.

⁽⁷⁾ The PS Clethodim rate should be 6 to 8 fl. oz./A for annual grass control in seedling alfalfa.

⁽⁸⁾ For the PS Clethodim plus LORSBAN tank mix, reduce the adjuvant rate down to 1.0 pt./A when the LORSBAN rate is 1.0 pt./A or higher.

Table 17. ROUNDUP READY VOLUNTEER CORN CONTROL IN ROUNDUP READY SOYBEAN WITH PS CLETHODIM HERBICIDE TANK MIX

| Roundup Ready Volunteer Corn Height (Inches) | PS Clethodim Rate fl. oz./A | Glyphosate ⁽¹⁾ rate for formulations with built in adjuvant | Adjuvant |
|----------------------------------------------|-----------------------------|-------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|
| <12 | 4 | 1.0 to 2.0 lb. ai./A (Approximately equivalent to 22 to 44 fl. oz./A of ROUNDUP Weather MAX) | Non-ionic surfactant @ 0.125 to 0.25% v/v plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals./ of carrier |
| 12 to 18 | 5 | | |
| 18 to 24 | 6 | | |

| Roundup Ready Volunteer Corn Height (Inches) | PS Clethodim Rate fl. oz./A | Glyphosate ⁽¹⁾ rate for formulations without built in adjuvant | Adjuvant |
|----------------------------------------------|-----------------------------|----------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|
| <12 | 4 | Up to 2.0 lb. ai./A (Equivalent to 32 to 64 fl. oz./A of ROUNDUP Original) | Crop oil concentrate @ 1 pt./A plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier |
| 12 to 18 | 5 | | |
| 18 to 24 | 6 | | |

⁽¹⁾ Glyphosate formulation must be labeled for use on Roundup Ready soybean.

THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label.
- Apply under favorable soil moisture and humidity which exist a few days after rainfall or within seven days after irrigation.
- Tank mix applications may sometimes result in reduced grass control. If regrowth occurs, or an additional flush of new grass emerges, make a second application of PS Clethodim, as specified in the respective size and rate tables.
- Do not tank mix PS Clethodim when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.
- This tank mix may be applied postemergence to ROUNDUP READY soybean up through the full flowering stage. Do not apply less than 60 days before harvest.
- Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants and trees, other than soybean with the ROUNDUP READY gene as severe injury or destruction will result.
- Do not allow the PS Clethodim plus ROUNDUP to mist, drip, drift or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur such as combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.

FALLOW LAND

DIRECTIONS FOR USE

PS Clethodim may be used to control annual and perennial grasses in land that has been left fallow the previous year and other non-producing agricultural areas. Apply PS Clethodim at 6 to 8 fl. oz./A for annual grasses and 8 to 16 fl. oz./A for perennial grasses. When both grass and broadleaf weeds are the target pest, PS Clethodim may be tank mixed with 2,4-D ester or BANVEL® SGF for broad spectrum control. When both annual and perennial grasses occur in the same field, use a minimum of 8 fl. oz./A PS Clethodim rate.

PRODUCT INFORMATION:

- Use a minimum spray volume of 5 gals./A for aerial applications and 15 gals./A for ground applications.
- Apply only to actively growing grasses when the first grass reaches the recommended weed height as specified by the Annual and Perennial Grasses section of this label.
- Annual grasses that emerge after the PS Clethodim application will not be controlled, and a second application may be necessary.
- The control of perennial grasses may require more than 1 application in non-tilled areas.
- Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.
- Do not apply to grasses that have tillered, formed seedheads or exceeded recommended growth stage.
- Do not use flood jet nozzles.
- Do not apply to drought stressed grasses.
- Do not mow area for 2 weeks prior to or after the PS Clethodim application.

Table 18. PS CLETHODIM IN TANK MIXES TO CONTROL ANNUAL AND PERENNIAL GRASSES IN FALLOW LAND

| PRODUCT | APPLICATION RATES/ACRE ⁽¹⁾ | | CROP OIL CONCENTRATE ⁽²⁾ (V/V) | |
|------------------------------------------------------|-----------------------------------------------------------------------------|-------------------|-------------------------------------------|-----|
| | ANNUAL GRASSES | PERENNIAL GRASSES | GROUND | AIR |
| PS Clethodim + 2,4-D ester or BANVEL SGF | 6 to 8 fl. oz. + 0.5 lb./A or See BANVEL SGF label for rates | 8 to 16 fl. oz. | 1% | 1% |

⁽¹⁾ Refer to PS Clethodim label for weed height and species control. Review BANVEL SGF and 2,4-D labels for crop restrictions, use rates and weeds controlled.

⁽²⁾ Always use a crop oil concentrate or methylated seed oil containing at least 15% emulsifier at the listed rate (but not less than 1 pt./A) in the finished spray volume.

| GRASS SUPPRESSION IN NON-CROP AREAS WITH PS CLETHODIM | | | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------|----------------------|-----------|
| GRASS SPECIES | WEED STAGE | RATE FL. OZ./ACRE | HIGH RATE |
| Annual and perennial grasses that exceed height claimed for control on height charts above. | Up to and including grasses in the seed head stage. | 12 | 16 |
| Do not apply as part of a tank mix when applying PS Clethodim for grass suppression. Add a crop oil concentrate at 1 qt./A by ground to the finished spray volume. | | | |

Table 19. PS CLETHODIM FOR THE CONTROL AND/OR SUPPRESSION OF TALL FESCUE IN NATIVE PRAIRIE WARM-SEASON GRASS RESTORATION PROJECTS

| PRODUCT | PRODUCT RATES | GRASS WEEDS CONTROLLED/SUPPRESSED | | WEED STAGES |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|--------------------------------------|----------------------------|--------------------------------------------|
| | | Common Name | Scientific Name | |
| PS Clethodim | 10 to 12 fl. oz./A | Tall Fescue | <i>Festuca arundinacea</i> | 4 to 6 inches tall (40 to 60% green-up) |
| Adjuvant: PS Clethodim must be applied with crop oil concentrate at 1 qt./A, plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A. Mixing Order: Thoroughly mix spray grade ammonium sulfate in water, add PS Clethodim, then add crop oil concentrate. | | | | |

SPECIAL APPLICATION INSTRUCTIONS/ RESTRICTIONS/ PRECAUTIONS

Burn or mow fields a minimum of 3 weeks prior to application to remove excess crop residue. Apply in the spring, at 40 to 60% tall fescue green-up, prior to emergence of warm-season grasses. Do not mow area for 2 weeks after the PS Clethodim application.

Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 PSI at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood jet nozzles.

Apply only to fields that have warm-season grasses established for 2 years. Applications of PS Clethodim to emerged warm-season grasses may cause injury. Do not apply to warm-season grasses grown for seed.

Do not graze treated fields or feed treated forage and/or hay to livestock. Do not plant any crop for 30 days after application, unless clethodim is registered for use in that crop.

NOTE: PS Clethodim applications are most effective if applied when average nighttime temperatures are consistently greater than or equal to 47°F.

Table 20. PS CLETHODIM FOR THE SUPPRESSION OF TALL FESCUE SEED-HEADS IN NON-PRODUCING AGRICULTURAL AREAS

| PRODUCT | PRODUCT RATE | SUPPRESSION | APPLICATION TIMING |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------|----------------------------------------------------------|----------------------------------|
| PS Clethodim | 1 1/2 to 2 fl. oz./A | Tall Fescue Seed-Heads (<i>Festuca arundinacea</i>) | (50 to 90% Tall Fescue green-up) |
| ADJUVANT: PS Clethodim must be applied with crop oil concentrate at 1 qt./A, plus a spray grade ammonium sulfate at 2.5 to 4 lb./A. Mixing Order: Thoroughly mix spray grade ammonium sulfate in water, add PS Clethodim Herbicide, then add crop oil concentrate. | | | |

SPECIAL APPLICATION INSTRUCTIONS/ RESTRICTIONS/ PRECAUTIONS

Apply at 50 to 90% tall fescue green-up.

Use the higher PS Clethodim rate if less tall fescue green matter is present.

Do not mow area for two weeks after the PS Clethodim application.

Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 psi at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood nozzles.

2,4-D ester may be added to this tank mix for broadleaf control (see 2,4-D ester label for weeds controlled).

Do not graze treated fields or feed treated forage and/or hay to livestock. Do not plant any crop for 30 days after application, unless clethodim is registered for use in that crop.

DIRECTIONS FOR USE IN ORNAMENTALS

For ornamental plant uses, PS Clethodim can be used to control labeled grass weeds in greenhouses, bathhouses, shadehouses, and around outdoor ornamentals, including nurseries, parks, roadside plantings, and structure landscapes.

IMPORTANT

PS Clethodim successfully controls weeds in newly transplanted and established non-grassy ornamentals. Plant tolerance to PS Clethodim at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of PS Clethodim have investigated the safety factor to ornamental plants not listed on the label.

The following plants have shown a tolerance for PS Clethodim applications:

ORNAMENTAL TREES

| COMMON NAME | SCIENTIFIC NAME |
|-----------------------|--------------------------------|
| Alder, Red | <i>Alnus rubra</i> |
| Ash | <i>Fraxinus</i> spp. |
| Basswood | <i>Tilia</i> spp. |
| Birch, European White | <i>Betula pendula</i> |
| Birch, River | <i>Betula nigra</i> |
| Birch, White | <i>Betula papyrifera</i> |
| Crabapple, Flowering | <i>Malus halliana</i> |
| Dogwood, Flowering | <i>Cornus florida</i> |
| Golden Chain Tree | <i>Laburnum anagyroides</i> |
| Maples | <i>Acer</i> spp. |
| Mulberry, White | <i>Morus alba</i> |
| Oaks | <i>Quercus</i> spp. |
| Olive, Wild | <i>Elaeagnus angustifolia</i> |
| Redbud, Eastern | <i>Cercis Canadensis</i> |
| Sweet Gum, American | <i>Liquidambar styraciflua</i> |

GROUND COVERS

| COMMON NAME | SCIENTIFIC NAME |
|--------------------|-------------------------------|
| Bugleweed, Carpet | <i>Ajuga reptans</i> |
| Ivy, English | <i>Hedera helix</i> |
| Japanese Spurge | <i>Pachysandra terminalis</i> |
| Lilyturf | <i>Liriope muscari</i> |
| Moneywort | <i>Lysimachia nummularia</i> |
| Mondo Grass, White | <i>Ophiopogon jaburan</i> |
| Mondo Grass Dwarf | <i>Ophiopogon japonicas</i> |
| Periwinkle, Lesser | <i>Vinca minor</i> |

GARDEN FLOWERS AND PLANTS

| COMMON NAME | SCIENTIFIC NAME |
|-------------------------|--------------------------------|
| Ageratum | <i>Ageratum</i> spp. |
| Alyssum*, Sweet | <i>Lobularia maritima</i> |
| Asparagus Fern | <i>Asparagus setaceus</i> |
| Bleeding Heart | <i>Dicentra spectabilis</i> |
| Cast Iron Plant | <i>Aspidistra elatior</i> |
| Chrysanthemum | <i>Chrysanthemum</i> spp. |
| Cinquefoil | <i>Potentilla</i> spp. |
| Coleus | <i>Coleus</i> spp. |
| Coralbells | <i>Heuchera sanguinea</i> |
| Cranesbill | <i>Geranium</i> spp. |
| Dahlia | <i>Dahlia</i> spp. |
| Daisy, Trailing African | <i>Osteospermum fruticosum</i> |
| Daylily | <i>Hemerocallis</i> spp. |
| Dusty Miller | <i>Senecio cineraria</i> |
| Euonymus | <i>Euonymus</i> spp. |
| Gazania | <i>Gazania</i> spp. |
| Geranium, House | <i>Pelargonium hortorum</i> |
| Heather, False | <i>Cuphea hyssopifolia</i> |

GARDEN FLOWERS AND PLANTS (cont'd.)

| COMMON NAME | SCIENTIFIC NAME |
|--------------------|------------------------------|
| Hosta | <i>Hosta fortunei</i> |
| Iris | <i>Iris</i> spp. |
| Jasmine Tobacco | <i>Nicotiana glauca</i> |
| Loosestrife | <i>Lythrum salicaria</i> |
| Marigold | <i>Tagetes</i> spp. |
| Partridgeberry | <i>Mitchella repens</i> |
| Petunia* | <i>Petunia hybrida</i> |
| Phlox | <i>Phlox</i> spp. |
| Pinks | <i>Dianthus</i> spp. |
| Portulaca | <i>Portulaca grandiflora</i> |
| Salvia | <i>Salvia</i> spp. |
| Saxifrage | <i>Saxifrage</i> spp. |
| Sedum | <i>Sedum</i> spp. |
| Selloum | <i>Philodendron selloum</i> |
| Snapdragon* | <i>Antirrhinum majus</i> |
| Sweet Flag | <i>Acorus gramineus</i> |
| Tickseed | <i>Coreopsis grandiflora</i> |
| Touch-Me-Not | <i>Impatiens</i> spp. |
| Verbena | <i>Verbena</i> spp. |
| Violet | <i>Viola</i> spp. |
| Yarrow, Common | <i>Achillea millefolium</i> |
| Zinnia | <i>Zinnia elegans</i> |

* Slight foliage or flower speckling has been observed on these species.

SHRUBS

| COMMON NAME | SCIENTIFIC NAME |
|---------------------------|------------------------------------|
| Abelia | <i>Abelia</i> spp. |
| Anise, Purple | <i>Illicium floridanum</i> |
| Aucuba | <i>Aucuba</i> spp. |
| Azalea* | <i>Rhododendron</i> spp. |
| Bamboo | <i>Bambusa</i> spp. |
| Barberry, Japanese | <i>Berberis thunbergii</i> |
| Barberry, Magellan | <i>Berberis buxifolia</i> |
| Bayberry | <i>Myrica pensylvanica</i> |
| Bottlebrush | <i>Callistemon citrinus</i> |
| Boxwood, Common | <i>Buxus sempervirens</i> |
| Camellia, Common | <i>Camellia japonica</i> |
| Candytuft | <i>Iberis sempervirens</i> |
| Cleyera | <i>Cleyera japonica</i> |
| Coralberry | <i>Ardisia crenata</i> |
| Crape Myrtle | <i>Lagerstroemia indica</i> |
| Coyote Brush | <i>Baccharis pilularis</i> |
| Fig, Creeping | <i>Ficus pumila</i> |
| Gardenia | <i>Gardenia</i> spp. |
| Holly | <i>Ilex</i> spp. |
| Honeysuckle | <i>Lonicera</i> spp. |
| Indian Hawthorn | <i>Raphiolepis indica</i> |
| Jasmine | <i>Jasminum</i> spp. |
| Jasmine, Asiatic | <i>Trachelospermum asiaticum</i> |
| Jasmine, Star | <i>Trachelospermum jasminoides</i> |
| Juniper | <i>Juniperus</i> spp. |
| Lantana | <i>Lantana</i> spp. |
| Nandina* Bamboo, Heavenly | <i>Nandinia domestica</i> |
| Oleander, Common | <i>Nerium oleander</i> |
| Oregon Grape | <i>Mahonia aquifolium</i> |

SHRUBS (cont'd.)

| COMMON NAME | SCIENTIFIC NAME |
|---------------------------|---------------------------|
| Photinia | <i>Photinia</i> spp. |
| Pittosporum | <i>Pittosporum</i> spp. |
| Podocarpus | <i>Podocarpus</i> spp. |
| Privet | <i>Ligustrum</i> spp. |
| Pyracantha | <i>Pyracantha</i> spp. |
| Rhododendron | <i>Rhododendron</i> spp. |
| Rose | <i>Spiraea bumalda</i> |
| Sweet Olive | <i>Osmanthus fragrans</i> |
| Viburnum | <i>Viburnum tinus</i> |
| Wisteria | <i>Wisteria</i> spp. |
| Yellow Sage/Shrub Verbena | <i>Lantana camara</i> |

* Slight foliage or flower speckling has been observed on these species.

ANNUAL GRASSES IN ORNAMENTALS

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass and weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

| GRASS SPECIES | SCIENTIFIC NAME | WEED* HEIGHT INCHES | RATE FL. OZ./ACRE ⁽¹⁾ | HIGH RATE ⁽²⁾ |
|------------------------|---------------------------------|------------------------|-------------------------------------|--------------------------|
| Barnyardgrass | <i>Echinochloa crusgalli</i> | 2 to 8 | 8 | 16 |
| Broadleaf Signalgrass | <i>Brachiaria platyphylla</i> | 2 to 6 | 8 | 16 |
| Brome | | | | |
| California | <i>Bromus carinatus</i> | 2 to 6 | 8 | 16 |
| Cheat | <i>Bromus secalinus</i> | 2 to 6 | 8 | 16 |
| Downy | <i>Bromus tectorum</i> | 2 to 6 | 8 | 16 |
| Ripgut | <i>Bromus diandrus</i> | 2 to 6 | 8 | 16 |
| Canarygrass | <i>Phalaris canariensis</i> | 1 to 4 | 8 | 16 |
| Crabgrass | | | | |
| Hairy | <i>Digitaria adscendens</i> | 2 to 6** | 8 | 16 |
| Large | <i>Digitaria sanguinalis</i> | 2 to 6** | 8 | 16 |
| Smooth | <i>Digitaria ischaemum</i> | 2 to 6** | 8 | 16 |
| Southern | <i>Digitaria ciliaris</i> | 2 to 6** | 8 | 16 |
| Crowfootgrass | <i>Dactyloctenium aegyptium</i> | 2 to 6** | 8 | 16 |
| Fall Panicum | <i>Panicum dichotomiflorum</i> | 2 to 8 | 8 | 16 |
| Field Sandbur | <i>Cenchrus incertus</i> | 2 to 6 | 8 | 16 |
| Foxtail | | | | |
| Giant | <i>Setaria faberi</i> | 2 to 12 | 8 | 16 |
| Green | <i>Setaria viridis</i> | 2 to 8 | 8 | 16 |
| Yellow | <i>Setaria glauca</i> | 2 to 8 | 8 | 16 |
| Goosegrass | <i>Eleusine indica</i> | 2 to 6** | 8 | 16 |
| Itchgrass | <i>Rottboellia cochlin</i> | 2 to 6 | 8 | 16 |
| Junglerice | <i>Echinochloa colona</i> | 2 to 6 | 8 | 16 |
| Lovegrass (Stinkgrass) | <i>Eragrostis ciliaris</i> | 2 to 6 | 8 | 16 |
| Rabbitsfootgrass | <i>Polygonon monspeliensis</i> | 1 to 4 | 8 | 16 |

ANNUAL GRASSES IN ORNAMENTALS (cont'd.)

| GRASS SPECIES | SCIENTIFIC NAME | WEED* HEIGHT INCHES | RATE FL. OZ./ACRE⁽¹⁾ | HIGH RATE⁽²⁾ |
|-------------------------|--------------------------------|--------------------------------|--------------------------------------------|--------------------------------|
| Red Rice | <i>Oryza sativa</i> | 1 to 3 | 8 | 16 |
| Ryegrass | | | | |
| Hardy | <i>Lolium remotum</i> | 2 to 6 | 8 | 16 |
| Italian | <i>Lolium multiflorum</i> | 2 to 6 | 8 | 16 |
| Seedling Johnsongrass | <i>Sorghum halepense</i> | 4 to 10 | 8 | 16 |
| Shattercane | <i>Sorghum bicolor</i> | 6 to 18 | 8 | 16 |
| Southwestern Cupgrass | <i>Eriochloa gracilis</i> | 2 to 6 | 8 | 16 |
| Sprangletop | | | | |
| Amazon | <i>Leptochloa panicoides</i> | 2 to 6 | 8 | 16 |
| Bearded | <i>Leptochloa fascicularis</i> | 2 to 6 | 8 | 16 |
| Mexican | <i>Leptochloa uninervia</i> | 2 to 6 | 8 | 16 |
| Red | <i>Leptochloa filiformis</i> | 2 to 6 | 8 | 16 |
| Texas Panicum | <i>Panicum texanum</i> | 2 to 6 | 8 | 16 |
| Volunteer Cereals | | | | |
| Barley | <i>Hordeum vulgare</i> | 2 to 6 | 8 | 16 |
| Oats | <i>Avena sativa</i> | 2 to 6 | 8 | 16 |
| Rye | <i>Secale cereale</i> | 2 to 6 | 8 | 16 |
| Wheat | <i>Triticum aestivum</i> | 2 to 6 | 8 | 16 |
| Volunteer Corn | <i>Zea mays</i> | 4 to 12 | 6 | 8 |
| Volunteer Corn | <i>Zea mays</i> | 12 to 24 | 8 | 16 |
| Volunteer Grain Sorghum | <i>Sorghum bicolor</i> | 8 to 12 | 8 | 16 |
| Wild Oats | <i>Avena fatua</i> | 2 to 6 | 8 | 16 |
| Wild Proso Millet | <i>Panicum miliaceum</i> | 2 to 10 | 8 | 16 |
| Witchgrass | <i>Panicum capillare</i> | 2 to 8 | 8 | 16 |
| Woolly Cupgrass | <i>Eriochloa villosa</i> | 2 to 8 | 8 | 16 |

* Generally occurs between 3-leaf stage and tillering.

** Length of lateral growth.

⁽¹⁾ 8 fl. oz./A = approximately 0.2 fl. oz./1000 sq. ft.

⁽²⁾ 16 fl. oz./A = approximately 0.4 fl. oz./1000 sq. ft.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

ANNUAL BLUEGRASS CONTROL WITH PS CLETHODIM IN ORNAMENTALS

| GRASS SPECIES | WEED STAGE | RATE FL. OZ./ACRE | HIGH RATE |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|------------------------------|------------------|
| Annual Bluegrass (<i>Poa annua</i>) | to 4-leaf | 6 | 16 |
| <p>Apply under favorable soil moisture and humidity that exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).</p> <p>Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.</p> <p>Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.</p> <p>Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).</p> | | | |

PERENNIAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

| GRASS SPECIES | WEED HEIGHT (inches) | RATE FL. OZ./ACRE ⁽¹⁾ | HIGH RATE ⁽²⁾ |
|-------------------------------------------------|-------------------------|-------------------------------------|--------------------------|
| <i>Bermudagrass (Cynodon dactylon)</i> | | | |
| First Application | 3 (or up to 6" runners) | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 3 (or up to 6" runners) | 8 | 16 |
| <i>Quackgrass (Elytrigia repens)</i> | | | |
| First Application | 4 to 8 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 4 to 8 | 8 | 16 |
| <i>Rhizome Johnsongrass (Sorghum halepense)</i> | | | |
| First Application | 12 to 24 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 6 to 18 | 6 | 8 |
| <i>Wirestem Muhly (Muhlenbergia frondosa)</i> | | | |
| First Application | 4 to 8 | 8 | 16 |
| Repeat Application(s) (if regrowth occurs) | 4 to 8 | 8 | 16 |

⁽¹⁾ 8 fl. oz./A = approximately 0.2 fl. oz./1000 sq. ft.

⁽²⁾ 16 fl. oz./A = approximately 0.4 fl. oz./1000 sq. ft.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals.
(0.25% v/v)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Open dumping is prohibited.

PESTICIDE STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place.

Do not store diluted spray.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Prime Source, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Prime Source, LLC and Seller harmless for any claims relating to such factors.

Prime Source, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Prime Source, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, PRIME SOURCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Prime Source, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PRIME SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Prime Source, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Prime Source, LLC.

Asana® is a registered trademark of DuPont.

Authority Broadleaf®, Canopy®, Classic® and Synchrony® are registered trademarks of E.I. DuPont de Nemours & Co. Inc.

STS™ is a trademark of E.I. DuPont de Nemours & Co. Inc.

Banvel®, IMI-CORN® and Storm® are registered trademarks of BASF Corporation.

Basagran® is a registered trademark of BASF Ag.

Baythroid® and Sencor® are registered trademarks of Bayer Corporation.

Betamix® and Betanex® are registered trademarks of The Aventis Group.

Blazer® is a registered trademark of United Phosphorus, Inc.

Bronate®, Bronate® Advanced® are registered trademarks of Bayer CropScience.

Buctril® and Turbo® are registered trademarks of Bayer Corporation.

Cobra® and Resource® are registered trademarks of Valent U.S.A. Corporation.

Danitol® is a registered trademark of Sumitomo Chemical Co., Ltd.

Dual®, Dual II®, Dual Magnum®, Flexstar® and Warrior® are registered trademarks of Syngenta Corporation.

Eminent® is a registered trademark of Sipcam Agro USA, Inc.

Firstrate®, Frontrow®, Lorsban® and Stinger® are registered trademarks of Dow AgroSciences LLC.

Liberty Link® is a registered trademark of The Aventis Group.

Orthene® is a registered trademark of OMS Investments, Inc.

Pounce®, Prowl®, Pursuit®, Pursuit DG®, Raptor® are registered trademarks of FMC Corporation.

RhonoX® is a registered trademark of Nufarm Americas Inc.

Roundup Ready® and Roundup Ultra® are registered trademarks of the Monsanto Company.

Valor® is a registered trademark of Valent U.S.A. Corporation.