

SODIUM MOLYBDATE

**FOR THE CORRECTIONS OF MOLYBDENUM
DEFICIENCIES IN CERTAIN PLANTS.**

Guaranteed Analysis: **By Wt.**
Molybdenum as (Mo) 39.0%
Derived from Sodium Molybdate Dihydrate

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin and eyes. Wash thoroughly after use.

Southern Agricultural Insecticides, Inc.

P.O. Box 218 P.O. Box 429 P.O. Box 85
Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607

F672 **Net Weight: 1 Pound (.45 kg)**

DIRECTIO Citrus


For use in full volume sprays
100 gallons of water and 5

Spray: Mix at the rate of
and spray to runoff.

Drench: Mix at the rate of
and apply to the substrate
tion per crop.
(Molybdenum deficiency
middle leaves of the plant
margin of the leaf. This ch
sis. Affected leaves may

Do not exceed

Buyer assumes all risks of
material not in strict acco
precautions given herew



SODIUM MOLYBDATE

**FOR THE CORRECTIONS OF MOLYBDENUM
DEFICIENCIES IN CERTAIN PLANTS.**

Guaranteed Analysis: **By Wt.**
Molybdenum as (Mo) 39.0%
Derived from Sodium Molybdate Dihydrate

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin and eyes. Wash thoroughly after use.

Southern Agricultural Insecticides, Inc.

P.O. Box 218 P.O. Box 429 P.O. Box 85
Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607

F672 **Net Weight: 1 Pound (.45 kg)**

DIRECTIO Citrus

For use in full volume sprays
100 gallons of water and 5

Spray: Mix at the rate of
and spray to runoff.

Drench: Mix at the rate of
and apply to the substrate
tion per crop.
(Molybdenum deficiency
middle leaves of the plant
margin of the leaf. This ch
sis. Affected leaves may

Do not exceed

Buyer assumes all risks of
material not in strict acco
precautions given herew

SODIUM MOLYBDATE

FOR THE CORRECTIONS OF MOLYBDENUM DEFICIENCIES IN CERTAIN PLANTS.

Guaranteed Analysis: **By Wt.**
Molybdenum as (Mo) 39.0%
Derived from Sodium Molybdate Dihydrate

KEEP OUT OF REACH OF CHILDREN

CAUTION:

May be harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin and eyes. Wash thoroughly after use.

DIRECTIONS FOR USE

Citrus & Tomatoes: For use in full volume sprayers mix at the rate of 1-2 ozs. per 100 gallons of water and spray to runoff.

Poinsettia: Spray: Mix at the rate of 1-2 ozs. per 100 gallons of water and spray to runoff.

Drench: Mix at the rate of 1-2 ozs. per 100 gallons of water and apply to the substrate. Make only one drench application per crop.

(Molybdenum deficiency of Poinsettias may show on the middle leaves of the plant as a chlorotic band around the margin of the leaf. This chlorosis quickly gives way to necrosis. Affected leaves may also be distorted.)

Do not exceed recommended rates.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

F672

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998