

SUBTILEX® NG

Biological Fungicide

ACTIVE INGREDIENT:

Bacillus subtilis, strain MBI 600* 9.9 %

OTHER INGREDIENTS:..... 90.1 %

TOTAL: 100.0%

* Contains not less than 5.5×10^{10} viable spores per gram

KEEP OUT OF REACH OF CHILDREN

CAUTION

For First Aid See Inside Panel

EPA Reg. No. 71840-8 EPA Est. No. 67064-IA-001

Becker Underwood, Inc.

801 Dayton Avenue

PO Box 667

Ames, IA 50010

Tel. 515-232-5907 • 800-232-5907

www.beckerunderwood.com

Net Weight: 2 ounces (56.7 grams)

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information on SUBTILEX® NG, call 1-800-232-5907, Monday through Friday, 8 a.m. to 5 p.m. CST. After 5 p.m. call your poison control center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

Mixers/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95 or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

Exception: if the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is:

- coveralls
- chemical-resistant gloves (made of any waterproof material)
- shoes plus socks

PRODUCT INFORMATION

SUBTILEX® NG contains bacteria that colonize developing root and shoot systems of plants, suppressing by competition disease organisms such as *Fusarium*, *Rhizoctonia*, , and *Pythium* as well as those organisms causing powdery mildew. Protection against root and soil borne pathogens is extended throughout the growing season as bacteria grow with the roots. As a result of this biological protection, vigorous root and shoot systems are established by treated plants, resulting in more uniform stands and greater yields.

SUBTILEX® NG is for use in soil or growing media, and for foliar applications to greenhouse-grown crops.

Apply SUBTILEX® NG using conventional application equipment as well as irrigation systems commonly used for chemigation.

FOR USE AS A FOLIAR TREATMENT OF GREENHOUSE CROPS

SUBTILEX® NG provides control of powdery mildew. SUBTILEX® NG is most effective as a preventative treatment. Apply when environmental conditions are favorable for disease development, but prior to disease onset. SUBTILEX® NG can be tank-mixed with most fungicides, insecticides, and fertilizers, but determine physical compatibility prior to use by mixing proportional quantities of the products in water.

Do not mix SUBTILEX® NG with any product containing a label prohibition against such mixing. When tank-mixing SUBTILEX® NG with any other registered foliar treatment products (insecticides, fungicides, fertilizers, etc.), observe the most restrictive of the labeling limitations and precautions of all products used in mixtures. Do not exceed label dosage rates.

APPLICATION RATES

For Greenhouse Crops: Apply SUBTILEX® NG at a rate of 0.4 – 1.2 oz/1000 ft² (11 – 33 g/100 m²) at 7 to 10 day intervals as needed. Use the stated higher rates of SUBTILEX® NG and the stated shorter application intervals when severe disease pressure is anticipated. Mix and apply SUBTILEX® NG in a sufficient volume of water to ensure uniform dispersion of product in the spray tank and thorough coverage of foliage and shoot tissue. Minimum application volume is 1 gallon per 1000 ft².

FOR USE AS A SOIL OR GROWING MEDIA TREATMENT

Apply SUBTILEX® NG as a water-based slurry to soil or growing media for preventative control and suppression of plant root pathogens *Rhizoctonia spp.*, *Pythium spp.* and *Fusarium spp.* SUBTILEX® NG can be tank-mixed with other registered insecticides, nematicides, fungicides or fertilizers. Prior to mixing, determine physical compatibility by mixing proportional quantities of the products in water.

Do not mix SUBTILEX® NG with any product containing a label prohibition against such mixing. When tank-mixing SUBTILEX® NG with any other soil or growing media treatment products, observe the most restrictive of the labeling limitations and precautions of all products used in mixtures. Do not exceed label application dosage rates.

APPLICATION RATES:

For pre-plant growing media amendment applications.

Apply SUBTILEX® NG at a rate of 0.05 – 0.07 oz/ cubic yard of soil or growing media (1.8 – 2.6 g/cubic meter). Use the higher rate when environmental conditions are favorable for disease development. Apply SUBTILEX® NG as a water-based slurry in a volume of water sufficient for uniform distribution. Typical application volume is 1 – 20 gal/cubic yard (5 – 100 L/cubic meter) of soil or growing media. Ensure product is thoroughly mixed into the soil or growing media.

For greenhouse post-plant applications.

Mix 0.2 – 0.4 oz. of SUBTILEX® NG in 100 gallons of water (15 – 30 g /1000 L). Use the higher rate when environmental conditions are favorable for disease development. Constant agitation is required to maintain SUBTILEX® NG in suspension. Apply evenly with conventional application equipment to thoroughly soak the growing media or soil through the root zone.

Conversion chart

Teaspoons		Ounces		Grams
0.38 tsp	=	0.05 oz	=	1.5
½ tsp	=	0.07 oz	=	2.0
1½ tsp	=	0.2 oz	=	5.7
3 tsp	=	0.4 oz	=	11.3

Container size	Min. drench volume (fl. oz. [mL])	Approximate number of containers treated per 100 gallons
Standard 4 inch (10 cm) round pot	1.5 [44]	8530
Standard 6 inch (15 cm) round pot	5.5 [163]	2330
Standard 8 inch (20 cm) round pot	12.75 [377]	1000

Begin applications during or after seeding, sticking of cuttings, or transplanting to pots, trays or containers, or when environmental conditions are favorable for disease development. For optimal control, use every 21-28 days throughout the growing cycle.

USES AND APPLICATION RATES FOR SELECTED CROPS

Greenhouse Crops

CROP	USE	DISEASES	RATE
Cucurbits: Cucumber, Cantaloupe, Melon, Muskmelon, Squash, Watermelon and other cucurbit crops	Soil or growing media	<i>Rhizoctonia</i> spp., <i>Pythium</i> spp. and <i>Fusarium</i> spp.	0.2-0.4 oz /100 gal* 15-30 g/1000 L*
	Foliar	Powdery mildew (<i>Sphaerotheca</i> spp., <i>Erysiphe</i> spp)	0.4-1.2 oz/1000 ft ² ** 11-33 g/100 m ² **
Flowers, Bedding Plants, Ornamentals, and Tropical Plants	Soil or growing media	<i>Rhizoctonia</i> spp., <i>Pythium</i> spp. and <i>Fusarium</i> spp.	0.2-0.4 oz /100 gal* 15-30 g/1000 L*
Fruiting Vegetables: Pepper, Tomato, Eggplant, and other fruiting vegetables	Soil or growing media	<i>Rhizoctonia</i> spp., <i>Pythium</i> spp. and <i>Fusarium</i> spp.	0.2-0.4 oz /100 gal* 15-30 g/1000 L*
	Foliar	Powdery mildew (<i>Leveillula taurica</i> , <i>Oidiopsis</i> <i>taurica</i> , <i>Sphaerotheca</i> spp., <i>Erysiphe</i> spp.)	0.4-1.2 oz/1000 ft ² ** 11-33 g/100 m ² **

* Thoroughly soak soil or growing media through root zone

** Minimum application volume is 1 gallon per 1000 ft².

CHEMIGATION:

General Requirements –

- 1) Apply this product only through a drip (trickle) system or sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- 2) Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- 3) If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 5) A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Requirements for Chemigation Systems Connected to Public Water Systems –

- 1) Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 2) Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 4) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 5) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7) Apply SUBTILEX® NG at the end of the water application, and in sufficient water for adequate coverage without excessive run off. Set the metering pump to the selected label use rate. Agitate the pesticide supply tank throughout the application of SUBTILEX® NG.
- 8) Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Drip (Trickle) Chemigation –

- 1) The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5) The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

- 7) Apply SUBTILEX® NG at the end of the water application, and in sufficient water for adequate coverage without excessive run off. Set the metering pump to the selected label use rate. Agitate the pesticide supply tank throughout the application of SUBTILEX® NG.

Specific Requirements for Sprinkler Chemigation –

- 1) The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5) The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7) Apply SUBTILEX® NG at the end of the water application, and in sufficient water for adequate coverage without excessive run off. Set the metering pump to the selected label use rate. Agitate the pesticide supply tank throughout the application of SUBTILEX® NG.
- 8) Do not apply when wind speed favors drift beyond the area intended for treatment.

Application Instructions –

- 1) Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues, may cause product to lose effectiveness or strength.
- 2) Determine the treatment rates as indicated in the directions for use and make proper dilutions.
- 3) Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry place until used. Do not store this product near food, feed, seed, fertilizers or other pesticides.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

NOTICE – READ CAREFULLY BEFORE USING
CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Becker Underwood, Inc. warrants that this product conforms to the specifications on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.

To the extent permitted by applicable law, Becker Underwood, Inc. **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use: It is impossible to eliminate all risks associated with use of this product. Lack of performance, injury, or other unintended consequences may result because of such factors as use of product contrary to strict label instructions and established safe practice, abnormal conditions (such as excessive rainfall, drought, and Acts of God), presence of other materials, use in combination with other materials, the manner of application, or other factors, all of which are beyond the control of Becker Underwood, Inc. or the seller. All such risks shall be assumed by the buyer.

Limitations of Remedies: To the extent permitted by applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Becker Underwood, Inc.'s election, one of the following: (1) replacement of the amount of the product used; or (2) refund of the purchase price paid for the product. Becker Underwood Inc. shall not be liable for losses or damages resulting from handling, storage, or use of this product contrary to label instructions unless Becker Underwood, Inc. is promptly notified of such loss and damage in writing. To the extent permitted by applicable law, Becker Underwood, Inc. shall not be liable for consequential or incidental damages or losses. Becker Underwood, Inc. neither assumes nor authorizes any person to assume for it, any other liability in connection with the sale, storage, use, or handling of this product other than expressly set forth herein.

Another quality biological product from:

Becker Underwood, Inc.

801 Dayton Avenue

PO Box 667

Ames, IA 50010

Tel. 515-232-5907 • 800-232-5907

www.beckerunderwood.com

SUBTILEX is a Registered Trademark of Becker Underwood, Inc.
Bacillus subtilis, strain MBI 600 is a product of Becker Underwood, Inc.