

This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC

SELECT™

INSECTICIDE

Sub-Label B (Pages 31-36): For Use on Ornamental Plants to Suppress Aphids, Thrips and Whiteflies and Control Mites and Leafminers

ACTIVE INGREDIENT:

Abamectin* (CAS No. 65195-56-4 and 65195-55-3) 2.0%

OTHER INGREDIENTS:..... 98.0%

Total 100.0%

*1 gal. contains 0.15 lb. abamectin

EPA Reg. No. 89442-20

GROUP	6	INSECTICIDE
--------------	----------	--------------------

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside for complete First Aid, Precautionary Statements, Directions for Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

Manufactured For:

Prime Source, LLC
4609 E. Boonville-New Harmony Road
Evansville, IN 47725



PRIME SOURCE

This product is manufactured by:
Prime Source, LLC
4609 E. Boonville-New Harmony
Road
Evansville, IN 47725




This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC

SELECT™

INSECTICIDE

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 a.m. to 4:30 p.m. Pacific Time or your poison control center at 1-800-222-1222 .	
NOTE TO PHYSICIAN Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical absorbents (e.g., activated charcoal). If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements. In severe cases, observation should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.	



This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC SELECT™

INSECTICIDE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING/AVISO. Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wear goggles, face shield or safety glasses.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category B on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical-resistant gloves, such as barrier laminate or butyl rubber ≥ 14 mils
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Attention: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



PRIME SOURCE



This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC SELECT™

INSECTICIDE

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber > 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.


Do not re-enter treated areas until sprays have dried.

PRODUCT INFORMATION

Read and follow all label directions. Applying Abamectin 0.15EC Select in a manner that is inconsistent with this label may result in decreased control of listed pests or damage to plants. This product is an emulsifiable concentrate and must be mixed with an adequate amount of water and applied as a foliar spray in order to achieve acceptable coverage.

Phytotoxicity to Abamectin 0.15EC Select has been tested on a broad variety of ornamentals. Not all combinations of this product with various mix partners (such as adjuvants, surfactants or other pesticides) or series of applications of this product and other additives have been evaluated, so a test application to a portion of the plant(s) is advised prior making a broader application, to assess the potential for damage to plants.

The following species have exhibited phytotoxic effects upon application of Abamectin 0.15EC Select; therefore, **do not** use this product on Ferns (e.g., *Adiantum* spp.) or Shasta Daisy (*Leucanthemum* spp.).



This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.


ABAMECTIN 0.15EC SELECT™

INSECTICIDE

Pests Suppressed	Directions
Aphids Thrips Whiteflies	<p>Aphids, thrips and whitefly will be killed through direct contact with Abamectin 0.15EC Select application solution.</p> <p>Apply this product when pests first appear in the immature, young stage of development. Repeat treatment at 7 day intervals for 2-3 weeks.</p> <p>After initial application sequence, use different product(s) with alternative mode(s) of action to this product¹ for a minimum of 2-3 weeks.</p>
<p>¹Refer to the Resistance Management section below.</p> <p>Restrictions</p> <ul style="list-style-type: none">To avoid developing resistance in <i>Liriomyza</i> leafminers and mites, do not apply this product for suppression of pests listed in this table on chrysanthemums, gerbera and roses. These plants are primary hosts of <i>Liriomyza</i> leafminers and mites. Extra applications of Abamectin 0.15EC Select to suppress pests listed in this table on chrysanthemums, gerbera and roses will increase selection pressure and may lead to greater tolerance of Abamectin 0.15EC Select on <i>Liriomyza</i> leafminers and mites. Application of this product on chrysanthemums, gerbera and roses should only be for control of leafminers and mites.	
Pests Controlled	Directions
Leafminers	<p>Apply this product as required for control of leafminers.</p> <p>Repeat applications may be made in order to maintain control at intervals of 7 days or as required.</p> <p>For resistance management of <i>Liriomyza</i> leafminer, apply Abamectin 0.15EC Select a maximum of three consecutive times. Following applications of this product, rotate to treatment with a product containing the active ingredient cyromazine for 3 consecutive applications maximum. Following treatment with cyromazine, switch back to treatment with Abamectin 0.15EC Select, or another product. Base variation between products with alternative modes of action on the generation period of the <i>Liriomyza</i> leafminer so that successive generations are not treated with the same product/mode of action. See the Resistance Management section below.</p>
Mites	<p>Apply this product when mites are first observed.</p> <p>Repeat application may be made as required in order to maintain control.</p>



PRIME SOURCE



This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC SELECT™

INSECTICIDE

Combining Abamectin 0.15EC Select with a horticultural spray oil may improve residual control of listed pests. Combine horticultural spray oil at a rate of 0.5-1% of spray application solution volume for the treatment of the following:

- Christmas trees
- Field-grown woody ornamentals
- Landscape plants

A repeat application may be made in order to maintain control. Wait 7 days between repeat applications.

In order to assess plant sensitivity to oils, make a spray application to a portion of the plants and then watch for 2 weeks, prior to making a broader application to the rest of the target area. The likelihood of plant injury following treatment with horticultural oil can be increased by excesses of temperature. Read and follow the label directions of the oil product and any other ingredients in the spray application mix.

Resistance Management


If strains of pests develop a tolerance to this product, the effectiveness of treatment with Abamectin 0.15EC Select can be reduced. It is important to implement a resistance management strategy with Abamectin 0.15EC Select in order to control labeled pests that will produce multiple generations in a plant or in a year. Resistance management measures to consider:

- Vary products with alternative modes of action.
- Avoid use of the same product or mode of action to treat subsequent generations of labeled pests.
- Correct use of products in accordance with product labels (correct use rates, application intervals etc.).
- Use of non-chemical pest management measures e.g., beneficial arthropods, plant rotation between vulnerable and less affected plants, beneficial cultural practices.

Refer to your State Cooperative Extension Service for more on resistance management strategies and implementation.

Restrictions

- **Do not** apply Abamectin 0.15EC Select through an irrigation system of any type.
- **Do not** apply Abamectin 0.15EC Select to any ornamentals via aerial application.
- **Do not** apply Abamectin 0.15EC Select in citrus nurseries.



This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC SELECT™

INSECTICIDE


CROP USE DIRECTIONS

For application in the following:

- shadehouse, greenhouse and field-grown ornamentals
- foliage plants
- Christmas trees
- other woody ornamentals

Apply Abamectin 0.15EC Select at a minimum use rate of 8 fl. oz./A and do not exceed 16 fl. oz./A. Abamectin 0.15EC Select must be combined with an adequate amount of water in order to ensure thorough and uniform plant coverage.

Use Rate of Abamectin 0.15EC Select	Pests Controlled/Suppressed	Directions
4 fl. oz./100 gals. water	Mites: Carmine Spider Mite Eriophyid Mites: Rust Bud Mites European Red Mite Southern Red Mite Spruce Spider Mite Tarsonemid Mites: Cyclamen Broad Mites Twospotted Spider Mite	Combine the listed use rate with 200-400 gallons of water per acre. If more than 400 gallons of water are to be applied per acre, use a maximum rate of 16 fl. oz. Abamectin 0.15EC Select per acre. If less than 200 gallons of water are to be applied per acre, use a minimum of 8 fl. oz. Abamectin 0.15EC Select per acre. For control of Tarsonemid Mites, newly developing tissue may require multiple treatments.
	8 fl. oz./100 gals. water	Boxwood Leafminer
	<i>Liriomyza</i> Leafminers	Combine the listed use rate with 100-200 gallons of water per acre. If more than 200 gallons of water are to be applied per acre, use a maximum rate of 16 fl. oz. Abamectin 0.15EC Select per acre. If less than 100 gallons of water are to be applied per acre, use a minimum rate of 8 fl. oz. Abamectin 0.15EC Select per acre. To maintain control, additional applications can be made at intervals of 7 days, or as needed.
	Aphids Thrips Whiteflies	In order to achieve suppression, young immature pests must come into contact with the spray application.



This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC SELECT™

INSECTICIDE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:


Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

Clean up minor spills or leaks immediately and follow directions and precautions on this label. During cleanup and disposal of wastes, avoid contamination of facilities and equipment.



PRIME SOURCE



This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package for complete label verbiage. This product may not yet be available or approved for sale or use in your area.

ABAMECTIN 0.15EC SELECT™

INSECTICIDE

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Prime Source, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Prime Source, LLC and Seller harmless for any claims relating to such factors.

Prime Source, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Prime Source, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, PRIME SOURCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Prime Source, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PRIME SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Prime Source, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Prime Source, LLC.

[04092015 v1]