

15-0-15

Soluble Fertilizer with 11% Calcium

	BY WT.
TOTAL NITROGEN (N)	15.00%
14.00% Nitrate Nitrogen (N)	
1.00% Ammoniacal Nitrogen (N)	
AVAILABLE PHOSPHATE (P₂O₅)	0.00%
AVAILABLE POTASH (K₂O)	15.00%
Calcium as Ca	11.00%
Manganese (Mn)	0.05%
0.05% Chelated Manganese as Mn	
Boron (B)	0.02%
Copper (Cu)	0.05%
0.05% Chelated Copper as Cu	
Iron (Fe)	0.10%
0.10% Chelated Iron as Fe	
Molybdenum (Mo)	0.015%
Zinc (Zn)	0.05%
0.003% Chelated Zinc as Zn	

Chlorine (Cl) - not more than 1.00%
PLANT NUTRIENTS DERIVED FROM: Calcium Nitrate, Potassium Nitrate, Ammonium Nitrate, Ferric Sodium EDTA, Zinc Disodium EDTA, Manganous Disodium EDTA, Cupric Disodium EDTA, Sodium Borate, Sodium Molybdate,
 EDTA - Ethylenediaminetetraacetate. **ARTIFICIALLY COLORED**
 Potential Basicity: 340 lb. Calcium Carbonate equivalent per ton.

GUARANTEED BY: Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34220

CAUTION: KEEP OUT OF REACH OF CHILDREN.

May be harmful if swallowed. Avoid breathing dust, and contact with skin, eyes and clothing. If ingested, give two glasses of water, induce vomiting and call a physician. If in eyes or on skin flush with water. If on clothing, launder before reuse. Wash thoroughly after handling.

USE PRECAUTIONS: This material is compatible with most insecticides and fungicides. Do not use in combination with Bordeaux, lime, highly alkaline materials. If compatibility is not known, test on small area before use.

Timing, method and rates of application, differences in plant varieties, growing conditions and other influencing factors are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607

F 672 J1 **Net Weight: 25 lb (11.34 kg)**

DIRECTIONS FOR USE

CONSTANT FEED: Where injector systems are used, the following table may be used to determine the amount of fertilizer to use in concentrate tank. When constant feeding, it is often desirable to leach potted and benched plants about once a week in order to avoid excessive fertilizer build up, particularly on the soil surface.

NOTE: Indicated rates are intended as a guide for initial applications. Experience with the particular crop and frequent soil testing are best guides to proper fertilization.

OUNCES BY WEIGHT PER GALLON OF CONCENTRATE			CONTINUOUS FEEDING ppm N			
FINAL CONC.	1:50	1:100	1:128	1:150	Crop Type	ppm
N in ppm	2.25	4.50	5.76	6.75	Bedding Plants	100-150
50 ppm	4.50	9.00	11.52	13.50	Woody Plants	50-100
100 ppm	6.75	13.50	17.28	20.25	Tropical Foliage	150-200
150 ppm	9.00	18.00	23.04	27.00	Chrysanthemums	250-300
200 ppm				36.00	Geraniums	250-300
					Plugs (all types)	50-125

100 ppm Nitrogen Solution of this product will contain 83 ppm Potassium and 73 ppm Calcium.

