SASOUTHERN 50AG.

# **RITIONAL SPRAY**



1 Tablespoonful per Gallon of Spray

FOR USE ON CITRUS, AVOCADOS, MANGOS AND OTHER TROPICAL FRUITS.

**Net Contents:** 1 Pint (473 ml)

Net Wt.: 1 lbs. 4.5 Ozs.



F672

P.O. Box 218

Southern Agricultural Insecticides, Inc. 218 Palmetto, FI 34220

### CITRUS NUTRITIONAL SPRAY

CITRUS NUTRITIONAL SPRA
GUARANTEED ANALYSIS:
1.00% Under Soluble Magnesium (Mg)
1.00% Water Soluble Magnesium (Mg)
4.10%
4.10%
4.10%
4.10%
5.20% Combined Sulfur (S)
1.20% Water Soluble Iron (Fe)
1.20% Water Soluble Iron (Fe)
1.20% Under Soluble Iron (Fe) 
 1.20% Water Soluble Iron (Fe)

 Manganese (Mn)
 1.20%

 1.20% Water Soluble Manganese (Mn)

 Zinc (Zn)
 1.70%

 1.70% Water Soluble Zinc (Zn)
 1.70%

 Derived from: Magnesium Sulfate, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate.
 Manganese Sulfate

## CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin and eyes. Avoid breathing spray mist. Do not take internally.

This product WILL SEVERELY STAIN concrete, masonry and painted surfaces. Be extremely careful not to allow spray or runoff to contact with these materials. Do not mix

with other fertilizers, fungicides or insecticides unless compatibility is known.

To questions concerning this product, please email us at info@southernag.com our consult our website at www.southernag.com.

Information regarding the contents and levels of metals in this product is available at http://www.aapfco.org/metals.htm

# DIRECTIONS FOR USE CITRUS, AVOCADOS, MANGOS, TROPICAL FRUITS & ORNAMENTALS

MANGOS, IRUPICAL FRUITS &
ORNAMENTALS
Citrus Nutritional Spray provides the minor elements
most needed by Citrus and Tropical Fruits.
This product will benefit the plant at any time when
deficiency symptoms appear. It is most effective when
applied between December and February up to the
time flowering is completed.
Apply spray to thoroughly cover to the point of run-off.
(preventative rate may also be used on ornamentals)
For prevention of deficiencies:
Use 1 tablespoonful per gallon of water.
1-2 applications per year are normally sufficient.
For correction of a known deficiency:
Use 2 tablespoonfuls per gallon of water and make
at least 2 applications at two week intervals.
On mature trees, the amount of spray required is
approximately 1 gallon for each foot of height plus 5
gallons, (20 foot tree = 25 gal. of spray)
NOTE: Deficiency symptoms in many plants can be an
indication of improper pH. Soil pH should be checked
and adjusted if necessary.

#### FIRST AID

IF SWALLOWED: Drink 2 glasses of water or milk and seek medical attention. EYE OR SKIN CON-TACT: Flush thoroughly with a large volume of water. If irritation persists seek medical attention.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given herewith.