



GLUFOSINATE GROUP 10 HERBICIDE



NULLIFY A/P

A non-selective herbicide for post emergence broadcast use on canola, corn, cotton, and soybean designated as LibertyLink®. NULLIFY A/P may be used for weed control in non LibertyLink® cotton when applied with a hooded sprayer in-crop. NULLIFY A/P may also be applied as a broadcast burndown application before planting or prior to emergence of any variety of canola, corn, cotton, soybean or sugar beet. NULLIFY A/P may be used for post emergence weed control in listed tree vine and berry crops. NULLIFY A/P may also be applied for potato vine desiccation.

ACTIVE INGREDIENT:

Glufosinate ammonium*	24.5%
Other Ingredients:	75.5%
Total:	100.0%

*Liquid soluble concentrate containing 2.34 pounds of active ingredient per U. S. gallon
EPA Reg. No. 42750-258-53883

(A) EPA Est. No. 88746-GA-01

(B) EPA Est. No. 53883-TX-002

First letter of lot# corresponds to EPA Est. # where product was produced

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label find someone to explain it to you in detail.)

See inside label booklet for additional

PRECAUTIONARY STATEMENTS and DIRECTIONS FOR USE

MANUFACTURED FOR: CONTROL SOLUTIONS, INC.

5903 Genoa-Red Bluff | Pasadena, TX 77507

EPA 080218

NET CONTENTS: 1 GALLON



FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN THE EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Do not give any liquid to the person.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE AT 1-800-424-9300.**

NOTE TO PHYSICIAN: If this product is ingested endotracheal intubation and gastric lavage should be performed as soon as possible followed by charcoal and sodium sulfate administration.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before use. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as Natural Rubber). Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants, shoes and socks

Applicators using groundboom equipment with open cabs to treat cotton must wear:

- Long sleeved shirt and long pants, shoes and socks plus chemical-resistant gloves.

Mixer/loaders supporting groundboom applications in corn, canola, soybeans, cotton, citrus fruit, pome fruit, stone fruit and olives must wear:

- Long sleeved shirt and long pants, shoes and socks plus chemical-resistant gloves.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating drinking chewing gum using tobacco or using the toilet
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d) (4 6)] the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present. Do not apply to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters or rinsate.

This pesticide is toxic to vascular plants and must be used strictly in accordance with the drift and run off precautions on this label in order to minimize off site exposures.

Under some conditions this product may have a potential to run off to surface water or adjacent land. Where possible use methods which reduce soil erosion such as no till limited till and contour plowing these methods also reduce pesticide run off. Use of vegetation filter strips along rivers creeks streams wetlands etc. or on the downhill side of fields where run off could occur to minimize water runoff is advised.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product until you have read the entire label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

In the State of New York Only: Not For Use in Nassau and Suffolk Counties.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses; and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 12 hours with the following exceptions:

- REI for workers engaged in scouting activities in corn, canola, and soybeans is 4 days.
- The REI for workers to move irrigation piping is 7 days for all crops.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls worn over short-sleeved shirt and short pants; chemical resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils; chemical resistant footwear plus socks; protective eyewear (goggles, face shield or safety glasses).

IMPORTANT CROP SAFETY INFORMATION READ BEFORE USING THIS PRODUCT

NULLIFY A/P may be applied as a **burndown treatment prior to planting or prior to emergence** of any variety of canola, corn, sweet corn*, cotton, olive, soybean or sugar beet. NULLIFY A/P may be applied to cotton not sensitive to the active ingredient in NULLIFY A/P using a hooded sprayer.

*Not for use in CA

Post emergence row crop applications of NULLIFY A/P may be made only to crops not sensitive to the active ingredient in this product. To the extent consistent with applicable law, Control Solutions, Inc. does not warrant the use of this product on crops other than those designated as LibertyLink[®] to safely withstand the application of NULLIFY A/P.

The basis of selectivity of NULLIFY A/P in crops is the presence of a gene in LibertyLink[®] crops which results in a plant that is not sensitive to the active ingredient in NULLIFY A/P. Crops not containing this gene will be sensitive to NULLIFY A/P, and severe crop injury and/or death may occur. Do not allow spray to contact foliage or green tissue of desirable vegetation other than crops not sensitive to the active ingredient in this product.

Applications to trees, vines and berries must avoid contact of NULLIFY A/P solution, spray, drift or mist with green bark, stems, or foliage, as injury may occur to trees, berries and vines. Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes or waxed containers. Contact of NULLIFY A/P with parts of trees, berries or vines other than mature brown bark can result in serious damage.

Mandatory Spray Drift Mitigation:

- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor blade vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- When applying to crops via aerial application equipment, applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- For aerial applications, do not release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is required for pilot safety.
- For ground applications and aerial applications, select nozzle and pressure that deliver medium to coarse spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with ASABE Standard 572.1.
- Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but do not exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.
- For non-crop vegetation management ground applications, apply with the nozzle height no more than 4 feet above the ground or target vegetation, unless necessitated by the application equipment. Examples would include roadside, railroad, utility rights of way, forestry and other industrial vegetation management applications where safety or natural barriers obstruct application.

Advisory Spray Drift Language

POLLINATOR ADVISORY STATEMENT:

This product contains an herbicide. Follow all label directions and precautions to minimize potential off-target exposure in order to prevent effects to non-target plants adjacent to the treated site which may serve as habitat or forage for pollinators.

Spray Drift Management:

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

Importance of Droplet Size:

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

Techniques for Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - Longer booms increase drift potential. Therefore, a shorter boom length is recommended.
- **Application Height** - Application more than 10ft. above the canopy increases the potential for spray drift.
- **Boom Height** - Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

Drift Reduction Technology (DRT)

The EPA Drift Reduction Technology (DRT) Program was developed to encourage the manufacture, marketing, and use of spray technologies scientifically verified to significantly reduce pesticide drift. The use of DRTs should result in significantly less pesticide from spray applications drifting and being deposited in areas not targeted by those applications, compared to spray technologies that do not meet the minimum DRT standard. EPA-verified drift reduction technologies (DRTs) and their ratings will be added to the following webpage as they become available: <https://www.epa.gov/reducing-pesticide-drift/epa-verified-and-rated-drift-reduction-technologies>

- **Wind** - Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.
Note: Local terrain can influence wind patterns. Every applicator needs to be familiar with local wind patterns and how they affect spray drift.
- **Temperature and Humidity** - When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.
- **Temperature Inversions** - Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if

fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

- **Shielded Sprayers** - Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

PRODUCT INFORMATION

NULLIFY A/P is a water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds in LibertyLink® canola, corn, cotton and soybean, and in trees, vines and berries. NULLIFY A/P may also be applied for potato vine desiccation. NULLIFY A/P may also be applied as a broadcast burndown application before planting or prior to emergence of any variety of canola, corn, sweet corn*, cotton, olive, soybean or sugar beet.

*Not for use in CA

NULLIFY A/P is only foliar-active with little or no activity in soil. Weeds that emerge after application will not be controlled. Apply NULLIFY A/P to actively growing weeds as described in the **Weed Control for Row Crops** section to get maximum weed control. Uniform, thorough spray coverage is necessary to achieve consistent weed control. Necrosis of leaves and young shoots occur within 2 to 4 days after application under good growing conditions.

- NULLIFY A/P is rainfast four (4) hours after application to most weed species; therefore, rainfall within four (4) hours may necessitate retreatment or may result in reduced weed control.
- Applications must be made between dawn and 2 hours before sunset to avoid the possibility of reduced lambsquarters palmer amaranth and velvetleaf control.
- Consult your local Cooperative Extension Service or Control Solutions, Inc. representative for guidelines on the optimum application timing for NULLIFY A/P in your region.
- Weed control may be reduced if application is made when heavy dew, fog and mist/rain are present; or when weeds are under stress due to environmental conditions such as drought, cool temperatures or extended periods of cloudiness.
- Warm temperatures, high humidity and bright sunlight improve the performance of NULLIFY A/P.
- NULLIFY A/P is a foliar-active material with little or no soil-residual activity.
- To maximize weed control, do not cultivate from 5 days before an application to 7 days after an application.

ROTATIONAL CROP RESTRICTIONS*

Rotational crop planting intervals following application of NULLIFY A/P are listed below. Failure to comply with these restrictions may result in illegal residues in rotated crops.

Rotational Crop	Plant Back Interval (Minimum Rotational Crop Planting Interval from Last Application)
Canola, Corn, Sweet Corn, Cotton, Rice, Soybean and Sugar Beet	May be planted at any time

Root and Tuber Vegetables, Leafy Vegetables, Brassica Leafy Vegetables and Small Grains (barley, buckwheat, oats, rye, teosinte, triticale, and wheat)	70 Days
All Other Crops	180 Days

*See Application Directions for Potato Vine Desiccation for Rotational Crop Restrictions specifically after NULLIFY A/P applications to potatoes.

WEED RESISTANCE MANAGEMENT

For resistance management, NULLIFY A/P Herbicide is a Group 10 herbicide (glutamine synthetase inhibitor). Any weed population may contain or develop plants naturally resistant to NULLIFY A/P and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

Contact your local sales representative, crop advisor or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of action for each target weed.

Fields should be scouted prior to application to identify the weed species present and their growth to determine if the intended application will be effective. Fields should be scouted after application to verify that the treatment was effective.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Control Solutions, Inc. representative. If resistance is suspected, treat weed escapes with an herbicide having a different mode of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

To delay herbicide resistance, take one or more of the following steps:

- **Start with clean fields.** Plant into weed-free fields and keep fields as weed free as possible. Effective tillage or the use of a burndown herbicide program can control emerged weeds prior to planting.
- **Scout fields.**
- **Diversified approach.** To the extent possible, use a diversified approach towards weed management. Whenever possible, incorporate multiple weed-control practices such as mechanical cultivation, biological management practices, and crop rotation.
- **Rotate crops.** Fields with difficult to control weeds should be rotated to crops that allow the use of herbicides with alternative modes of action or different management practices.
- **Control weed escapes.** To the extent possible, do not allow weed escapes to produce seeds, roots or tubers. Manage weeds at harvest and post-harvest to prevent a buildup of the weed seed-bank.
- **Clean equipment.** Prevent field-to-field and within-field movement of weed seed or vegetative propagules. Thoroughly clean plant residues from equipment before leaving fields.

- **Manage borders.** Prevent an influx of weeds into the field by managing borders.
- **Know your weeds, know your fields.** Identify weeds present in the field through scouting and field history and understand their biology. The weed-control program should consider all of the weeds present.
- **Rotate mechanisms of action.** Difficult to control weeds may require sequential applications of herbicides with differing mechanisms of action. Use a broad-spectrum soil-applied herbicide with a mechanism of action that differs from this product as a foundation in a weed-control program. Do not use more than two applications of this or any other herbicide with the same mechanism of action within a single growing season unless mixed with an herbicide with another mechanism of action with an overlapping spectrum for the difficult-to-control weeds. If resistance is suspected, treat weed escapes with an herbicide with a different MOA or use non-chemical methods to remove escapes.
- **Apply herbicide correctly.** Apply this herbicide at the correct timing and rate to control the most difficult weed in the field.

Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes. For more information on Weed Resistance Management, visit the Herbicide Resistance Action Committee (HRAC) on the web at <http://www.hracglobal.com>.

WEED CONTROL FOR ROW CROPS

Rates in ounces of formulated product per acre for the control of weeds as shown in the weed control tables. In weed populations with mixed species, apply at a rate needed for the species targeting less than three inch weeds.

Broadleaf Weeds Controlled (including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)			
Common Name	Scientific Name	22 fl oz/A (0.40 lbs ai/A)	29 - 43 fl oz/A (0.53 - 0.79 lbs ai/A)
		C = Control NR = Not Recommended S = Suppression	C = Control NR = Not Recommended S = Suppression
Amaranth, Palmer	<i>Amaranthus palmeri</i>	NR	C
Anoda, spurred	<i>Anoda cristata</i>	C	C
Beggarweed, Florida	<i>Desmodium tortuosum</i>	C	C
Black medic	<i>Medicago lupulina</i> L.	C	C
Blueweed, Texas	<i>Helianthus ciliaris</i> DC.	C	C
Buckwheat, wild	<i>Polygonum convolvulus</i>	C	C
Buffalobur	<i>Solanum cornutum</i>	C	C
Burcucumber	<i>Sicyos angulatus</i>	C	C
Canola, volunteer ¹	<i>Brassica spp.</i>	C ₁	C ₁

**Broadleaf Weeds Controlled
(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22 fl oz/A (0.40 lbs ai/A)	29 - 43 fl oz/A (0.53 - 0.79 lbs ai/A)
		C = Control NR = Not Recommended S = Suppression	C = Control NR = Not Recommended S = Suppression
Catchweed bedstraw (cleavers)	<i>Galium aparine</i> L.	C	C
Carpetweed	<i>Mollugo verticillata</i>	C	C
Chickweed, common	<i>Stellaria media</i>	C	C
Cocklebur, common	<i>Xanthium strumarium</i>	C	C
Copperleaf, hophornbeam	<i>Acalypha ostryaefolia</i>	C	C
Cotton, volunteer ¹	<i>Gossypium sp.</i>	C ₁	C ₁
Croton, tropic	<i>Croton glandulosus</i>	C	C
Croton, woolly	<i>Croton capitatus</i>	C	C
Eclipta	<i>Eclipta alba</i>	C	C
Devil's claw	<i>Proboscidea Louisiana</i>	C	C
Fleabane, annual	<i>Erigeron annuus</i>	C	C
Galinsoga, hairy	<i>Galinsoga ciliata</i>	C	C
Galinsoga, small flower	<i>Galinsoga parviflora</i>	C	C
Groundcherry, cutleaf	<i>Physalis angulate</i>	C	C
Geranium, cutleaf	<i>Geranium dissectum</i> L.	C	C
Hempnettle	<i>Galeopsis sp.</i>	C	C
Horsenettle, Carolina ²	<i>Solanum carolinense</i>	C ₂	C ₂
Jimsonweed	<i>Datura stramonium</i>	C	C
Knotweed	<i>Polygonum spec.</i>	C	C
Ladysthumb	<i>Polygonum persicaria</i>	C	C
Kochia	<i>Kochia scoparia</i>	C	C
Lambsquarters, common	<i>Chenopodium album</i>	C	C
Mallow, common	<i>Malva spec.</i>	C	C

Broadleaf Weeds Controlled
(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)

Common Name	Scientific Name	22 fl oz/A (0.40 lbs ai/A)	29 - 43 fl oz/A (0.53 - 0.79 lbs ai/A)
		C = Control NR = Not Recommended S = Suppression	C = Control NR = Not Recommended S = Suppression
Mallow, Venice	<i>Hibiscus trionum</i>	C	C
Marestail ³	<i>Coryza Canadensis</i>	S	C
Marsh-elder, annual	<i>Iva annua</i>	C	C
Morningglory, entireleaf	<i>Ipomoea hederacea</i> var. <i>intergriuscula</i>	C	C
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	C	C
Morningglory, pitted	<i>Ipomoea lacunose</i>	C	C
Morningglory, sharppod	<i>Ipomoea cordatotriloba</i>	C	C
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>	C	C
Morningglory, tall	<i>Ipomoea purpurea</i>	C	C
Mustard, wild	<i>Sinapis arvensis</i>	C	C
Nightshade, black	<i>Solanum nigrum</i>	C	C
Nightshade, eastern black	<i>Solanum ptycanthum</i>	C	C
Nightshade, hairy	<i>Solanum sarracoides</i>	C	C
Pennycress	<i>Thlaspi arvense</i>	C	C
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C	C
Pigweed, prostrate	<i>Amaranthus blitoides</i>	C	C
Pigweed, spiny	<i>Amaranthus spinosus</i>	C	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C	C
Pigweed, tumble	<i>Amaranthus albus</i>	C	C
Puncturevine	<i>Tribulus terrestris</i>	C	C
Purslane, common	<i>Portulaca oleracea</i>	C	C
Pusley, Florida	<i>Richardia scabra</i>	S	C
Ragweed, common	<i>Ambrosia artemisiifolia</i>	C	C

**Broadleaf Weeds Controlled
(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22 fl oz/A (0.40 lbs ai/A)	29 - 43 fl oz/A (0.53 - 0.79 lbs ai/A)
		C = Control NR = Not Recommended S = Suppression	C = Control NR = Not Recommended S = Suppression
Ragweed, giant	<i>Ambrosia trifida</i>	C	C
Senna coffee	<i>Cassia occidentalis</i>	C	C
Sesbania, hemp	<i>Sesbania herbacea</i>	C	C
Shepherd's-Purse	<i>Capsella bursa-pastoris</i>	C	C
Sicklepod (java bean)	<i>Senna obtusifolia</i>	C	C
Sida, prickly	<i>Sida spinosa</i> L.	C	C
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	C	C
Smell melon	<i>Cucumis melo</i> L. var. <i>Dudaim</i>	C	C
Sowthistle, annual	<i>Sonchus oleraceus</i> L.	C	C
Soybeans, volunteer ¹	<i>Glycine max</i>	C ₁	C ₁
Spurge, prostrate	<i>Euphorbia humifusa</i>	C	C
Spurge, spotted	<i>Euphorbia maculate</i> L.	C	C
Starbur, bristly	<i>Acanthospermum hispidum</i>	C	C
Sunflower, common	<i>Helianthus annuus</i>	C	C
Sunflower, prairie	<i>Corythucha pura</i>	C	C
Sunflower, volunteer	<i>Girassol</i>	C	C
Thistle, Russian ²	<i>Salsola kali</i>	S ₂	C ₂
Velvetleaf	<i>Abutilon theophrasti</i>	C	C
Waterhemp, common	<i>Amaranthus rudis</i>	NR	C
Waterhemp, tall	<i>Amaranthus tuberculatos</i>	NR	C

¹ Volunteer LibertyLink crops from the previous season will not be controlled.

² May require sequential applications for control.

³ For optimum control apply NULLIFY A/P on 6" mareetail.

**Grass Weeds Controlled
(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22 fl oz/A (0.40 lbs ai/A)	29 - 43 fl oz/A (0.53 - 0.79 lbs ai/A)
		C = Control NR = Not Recommended S = Suppression	C = Control NR = Not Recommended S = Suppression
Barley, volunteer ³	<i>Hordeum vulgare</i>	C ₃	C ₃
Barnyardgrass	<i>Echinochloa spec.</i>	C	C
Bluegrass, annual	<i>Poa annua L.</i>	C	C
Corn, volunteer	<i>Zea mays L.</i>	C ₁	C ₁
Crabgrass, large	<i>Digitaria sanguinalis</i>	C ₂	C ₂
Crabgrass, smooth ²	<i>Digitaria ischaemum</i>	C ₂	C ₂
Cupgrass, woolly	<i>Eriochloa villosa</i>	C	C
Foxtail, bristly	<i>Setaria verticillata</i>	C	C
Foxtail, giant	<i>Setaria faberi</i>	C	C
Foxtail, green	<i>Setaria viridis</i>	C	C
Foxtail, robust purple	<i>Setaria viridis</i>	C	C
Foxtail, yellow ²	<i>Pennisetum glaucum</i>	C ₂	C ₂
Goosegrass ³	<i>Eleusine indica</i>	C ₃	C ₃
Johnsongrass, seedling	<i>Sorghum halepense</i>	C	C
Junglerice	<i>Echinochloa colonum</i>	C	C
Millet, wild-proso	<i>Panicum miliaceum L.</i>	C	C
Millet, proso volunteer	<i>Milium vernale</i>	C	C
Oat, wild ²	<i>Avena fatua</i>	C	C
Panicum, fall	<i>Panicum dichotomiflorum</i>	C	C
Panicum, Texas	<i>Panicum texanum</i>	C	C
Rice, red	<i>Oryza sativa L.</i>	C	C
Rice, volunteer ¹	<i>Oryza sativa</i>	C ₁	C ₁
Sandbur, field ²	<i>Cenchrus pauciflorus</i>	S ₂	C ₂
Shattercane	<i>Sorghum vulgare PERS.</i>	C	C

Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>	C	C
Sprangletop	<i>Leptochloa spec.</i>	C	C
Sorghum, volunteer	<i>Sorghum sp.</i>	C	C
Stinkgrass	<i>Eragrostis ciliaris</i>	C	C
Wheat, volunteer ²	<i>Triticum spec.</i>	C ₂	C ₂
Witchgrass	<i>Panicum virgatum L.</i>	C	C

¹ Volunteer LibertyLink crops from the previous season will not be controlled. A timely cultivation 7 to 10 days after an application and/or retreatment 1021 days after the first application is advised for controlling dense clumps of volunteer corn or rice.

² For best control of yellow foxtail, field sandbur, crabgrass, and wild oats, treat prior to tiller initiation.

³ A sequential application may be necessary for control.

Biennial and Perennial Weeds Controlled (including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin Resistant Biotypes)		
For control of the biennial and perennial weeds listed below, tank mix partners or sequential applications NULLIFY A/P are advised by crop (see crop sections)		
Common Name	Scientific Name	29 - 43 fl oz/A (0.53 - 0.79 lbs ai/A)
		C = Control NR = Not Recommended S = Suppression
Alfalfa	<i>Medicago sativa L.</i>	C
Bermudagrass	<i>Cynodon dactylon</i>	C
Bindweed, field	<i>Convolvulus arvensis L.</i>	C
Bindweed, hedge	<i>Calystegia sepium</i>	C
Bluegrass, Kentucky	<i>Poa pratensis L.</i>	C
Blueweed, Texas	<i>Helianthus ciliaris DC.</i>	C
Bromegrass, smooth	<i>Bromus inermis</i>	C
Burdock	<i>Arctium sp.</i>	C
Bursage, woolyleaf	<i>Ambrosia grayi</i>	C
Chickweed, Mouse-ear	<i>Cerastium vulgatum L.</i>	C
Clover, red	<i>Trifolium pretense L.</i>	C
Dandelion	<i>Taraxacum officinale</i>	S
Dock, smooth	<i>Rumex spec.</i>	S
Dogbane, hemp	<i>Apocynum cannabinum</i>	C

**Biennial and Perennial Weeds Controlled
(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin Resistant Biotypes)**

For control of the biennial and perennial weeds listed below, tank mix partners or sequential applications NULLIFY A/P are advised by crop (see crop sections)

Common Name	Scientific Name	29 - 43 fl oz/A (0.53 - 0.79 lbs ai/A)
		C = Control NR = Not Recommended S = Suppression
Goldenrod, gray	<i>Solidago nemoralis</i>	S
Johnsongrass, rhizome	<i>Sorghum halepense</i>	S
Milkweed, common	<i>Asclepias syriaca</i>	S
Milkweed, honeyvine	<i>Ampelamus albidus</i>	C
Muhly, wirestem	<i>Muhlenbergia frondosa</i>	S
Nightshade, sliverleaf	<i>Solanum elaeagnifolium</i>	S
Nutsedge, purple	<i>Cyperus rotundus</i>	C
Nutsedge, yellow	<i>Cyperus ferax</i>	C
Orchardgrass	<i>Dactylis glomerata</i> L.	C
Poinsettia, wild	<i>Euphorbia heterophylla</i> L.	S
Pokeweed	<i>Phytolaccaceae</i>	C
Quackgrass	<i>Agropyron repens</i>	C
Sowthistle, perennial	<i>Sonchus arvensis</i> L.	C
Thistle, bull	<i>Cirsium vulgare</i>	S
Thistle, Canada	<i>Cirsium arvense</i>	C
Timothy	<i>Phleum pretense</i> L.	S
Wormwood, biennial	<i>Artemisia biennis</i>	C

APPLICATION AND MIXING PROCEDURES

Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment. Uniform thorough spray coverage is important to achieve consistent weed control.

Ground Application: Refer to the Rate Tables for proper application rates.

Aerial Application: Poor coverage will result in reduced weed control. See the Spray Drift Management section of this label for additional information on proper application of NULLIFY A/P.

COMPATIBILITY TESTING

If NULLIFY A/P is to be mixed with pesticide products not listed on this label, test the compatibility of the intended tank mixture prior to mixing the products in the spray tank. The following procedure assumes a spray volume of 25 gallons per acre. For other spray volumes, adjust the amount of the water used accordingly.

Check compatibility as follows:

1. Place 1.0 pint of water from the source that will be used to prepare the spray solution in a clear 1-quart jar.
2. For each pound of a dry tank mix partner to be applied per acre, add 1.5 teaspoons to the jar.
3. For each 16 fl oz of a liquid tank mix partner to be applied per acre, add 0.5 teaspoon to the jar.
4. For each 16 fl oz of NULLIFY A/P to be applied per acre, add 0.5 teaspoon to the jar.
5. After adding all the ingredients, place a lid on the jar and tighten. Invert 10 times to mix.
6. Let the mixture stand for 15 minutes, and evaluate the solution for uniformity and stability. Look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. If the tank mix partners are not compatible, do not use the mixture in a spray tank.
7. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the Storage and Disposal section of this label.

MIXING INSTRUCTIONS

Tank Mix: NULLIFY A/P may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. NULLIFY A/P cannot be mixed with any product containing a label prohibition against such mixing. Refer to the specific crop section for rate directions and other restrictions.

NULLIFY A/P must be applied with properly calibrated and clean equipment. NULLIFY A/P is formulated to mix readily in water. Prior to adding NULLIFY A/P to the spray tank, ensure that the spray tank is thoroughly clean, particularly if a herbicide with the potential to injure crops was previously used (see Cleaning Instructions).

Mix NULLIFY A/P with water to make a finished spray solution as follows:

1. Fill the spray tank half full with water.
2. Start agitation.
3. If mixing with a flowable/wettable powder tank mix partner. Prepare a slurry of the proper amount of the product in a small amount of water. Add the slurry to the spray tank.
4. Add the appropriate amount of ammonium sulfate (AMS) to the spray tank.
5. If mixing with a liquid tank mix partner, add the liquid mix partner next.
6. Complete filling the spray tank with water.
7. Add the proper amount of NULLIFY A/P and continue agitation.
8. If foaming occurs, use a silicone-based anti-foam agent.

Ensure that all spray system lines including pipes, booms, etc. have the correct concentration of spray solution by flushing out the spray system lines before starting the crop application.

If tank mix partners specified on this label are added, maintain good agitation at all times until contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep bypass line on or near bottom of tank to minimize foaming. Screen size in nozzles or line strainers must be 50 mesh or larger.

CLEANING INSTRUCTIONS

Before using NULLIFY A/P, thoroughly clean bulk storage tank, refillable tank, nurse tanks, spray tank, lines, and filter, particularly if a herbicide with the potential to injure crops was previously used. Equipment must be thoroughly rinsed using a commercial tank cleaner.

After using NULLIFY A/P, triple rinse the spray equipment and clean with a commercial tank cleaner before using for crops not labeled LibertyLink®. Make sure any rinsate or foam is thoroughly removed from spray tank and boom. Rinsate may be disposed following the pesticide disposal directions on this label.

APPLICATION DIRECTIONS FOR BURNDOWN USE

NULLIFY A/P may be applied as a **burndown treatment prior to planting or prior to emergence** of any variety of canola, corn, sweet corn, cotton, soybean, or sugar beet.

APPLICATION TIMING

Apply to small and actively growing weeds, targeting weeds less than 3 inches in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed control may be reduced when applications are made when heavy dew, fog and mist/rain are present or when weeds are under stress due to drought, cool temperatures or extended periods of cloudiness. NULLIFY A/P is a foliar-active material with little or no soil-residual activity. NULLIFY A/P is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment. For best results on lambsquarters, Palmer amaranth and velvetleaf control, make applications of Glufosinate between dawn and 2 hours before sunset.

APPLICATION RATES

Apply 29 to 43 fl oz/A (*0.53 to 0.79 lbs ai/A*) depending on crop, weed species and intention of post application use. Please see application charts below.

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

- **Cotton**, if environmental conditions prevent timely applications, a single application may be made of up to 43 fluid ounces per acre (*0.79 lb ai/A*) of NULLIFY A/P. **If more than 29 fluid ounces per acre (0.53 lb ai/A) are used in any single application, the annual total may not exceed 72 fluid ounces per acre (1.32 lbs ai/A), including all application timings.**
- In **canola, corn (sweet and field) and soybean**, if environmental conditions prevent timely applications, a single application may be made of up to 43 fluid ounces per acre (*0.79 lb ai/A*) of NULLIFY A/P. The year total may not exceed 43.0 fluid ounces per acre (*0.79 lb ai/A*) including all application timings.
- In **sugar beets**, if environmental conditions prevent timely applications, a single application may be made of up to 36 fluid ounces per acre (*0.66 lb ai/A*) of NULLIFY A/P. No additional applications of NULLIFY A/P may be made post emergence to the crop during the year.

ADJUVANTS

Ammonium sulfate (AMS) can be used at 1.5 lb/A to 3 lb/A. Rates are dependent on tankmix partners, environmental conditions, temperatures and potential for leaf burn. AMS has shown to improve weed control of difficult-to-control weeds like velvetleaf and lambsquarters, under difficult environmental conditions (low relative humidity) or hard water. An anti-foam agent is advised.

SURFACTANTS/OILS

The use of surfactants may be included. Please refer to the surfactant label for more detailed information.

NOZZLE SPRAY QUALITY

Use Medium to Coarse nozzles. NULLIFY A/P is a contact herbicide and requires proper nozzles with uniform thorough spray coverage to achieve optimum weed control. See nozzle section for more detailed information.

APPLICATION DIRECTIONS FOR CROPS

Crop	Burndown	In-Season Applications	Yearly Maximum
Canola, Soybean, Sweet Corn, Field Corn	29 - 43 fl oz/A (0.53 - 0.79 lb ai/A)	None	43 fl oz/A (0.53 lb ai/A)
Sugar beets	29 - 36 fl oz/A (0.53 - 0.66 lb ai/A)	None	36 fl oz/A (0.66 lb ai/A)
Cotton Use Pattern 1	29 fl oz/A (0.53 lb ai/A)	2 applications at 29 fl oz/A* (0.53 lb ai/A) Make second application 10-14 days after the first application.	87 fl oz/A (1.59 lbs ai/A)
Cotton Use Pattern 2	32 - 43 fl oz/A (0.58 - 0.79 lb ai/A)	1 application at 29 fl oz/A* (0.53 lb ai/A)	72 fl oz/A (1.32 lbs ai/A)

*Cotton containing the LibertyLink® trait OR with hooded sprayer for all varieties (see **COTTON** use directions).

APPLICATION DIRECTIONS FOR CROPS CONTAINING LIBERTYLINK® TRAIT

Crop	Burndown	In-Season Applications (LibertyLink® Varieties Only)	Yearly Maximum
Soybean, Field Corn	29 - 43 fl oz/A (0.53 - 0.79 lb ai/A)	1 to 2 applications at 29 - 43 fl oz/A (0.53 - 0.79 lb ai/A) For soybeans, make second application at least 5 days after the first application. For field corn, make second application at least 7 days after first application.	87 fl oz/A (1.59 lbs ai/A)
Sweet Corn	22 fl oz/A (0.40 lb ai/A)	1 to 2 applications at 22 fl oz/A (0.40 lb ai/A) Make second application at least 7 days after the first application.	44 fl oz/A (0.80 lb ai/A)

Crop	Burndown	In-Season Applications (LibertyLink® Varieties Only)	Yearly Maximum
Canola	29 - 43 fl oz/A (0.53 - 0.79 lb ai/A)	1 to 2 applications at 29 fl oz/A (0.53 ai/A) Make second application at least 10 days after the first application	87 fl oz/A (1.59 lbs ai/A)
Cotton Use Pattern 1	29 fl oz/A (0.53 lb ai/A)	1 to 2 applications at 29 fl oz/A (0.53 lb ai/A) Make second application 10-14 days after the first application.	87 fl oz/A (1.59 lb ai/A)
Cotton Use Pattern 2	30 - 43 fl oz/A (0.55 - 0.79 lb ai/A)	1 application at 29 fl oz/A (0.53 lb ai/A)	72 fl oz/A (1.32 lbs ai/A)
Sugar Beets	29 - 36 fl oz (0.53 - 0.66 lb ai/A)	1 application at 29 fl oz/A (0.53 lb ai/A)	60 fl oz/A (1.10 lbs ai/A)

APPLICATION DIRECTIONS FOR USE ON SUGAR BEETS

Apply NULLIFY A/P only to sugar beets labeled as LibertyLink. NULLIFY A/P is a contact herbicide and requires uniform, thorough spray coverage to achieve optimum weed control.

APPLICATION TIMING

Applications of NULLIFY A/P on sugar beets containing the LibertyLink trait may be made from the cotyledon stage up to the 10-leaf stage of the sugar beet.

Apply to small and actively growing weeds, targeting weeds less than 3 inches in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed control may be reduced when applications are made when heavy dew, fog and mist/rain are present or when weeds are under stress due to drought, cool temperatures or extended periods of cloudiness. NULLIFY A/P is a foliar-active material with little or no soil-residual activity. NULLIFY A/P is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment. For best result, on lambsquarters, Palmer amaranth and velvetleaf control, make applications of Glufosinate between dawn and 2 hours before sunset.

APPLICATION RATES

Apply 29 - 36 fluid ounces per acre (0.53 - 0.66 lb ai/A) depending on weed species, size and density per weed chart. If a second application is needed, make the second application in a minimum of 10 days after the first application. The maximum annual rate of NULLIFY A/P on sugar beets is 60 fl oz/A (1.10 lbs ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

ADJUVANTS

Ammonium sulfate (AMS) may be used at 1.5 to 3.5 lb/A. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (such as temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds like lambsquarters and velvetleaf under difficult environmental conditions (such as low relative humidity) or hard water. The use of an anti-foam agent is advised.

SURFACTANTS/OILS

The use of additional surfactants or crop oils in tank mixes with NULLIFY A/P may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

NOZZLE SPRAY QUALITY

Use medium to coarse nozzles. NULLIFY A/P is a contact herbicide and requires proper nozzles with uniform thorough spray coverage to achieve optimum weed control.

See **SPRAY DRIFT MANAGEMENT** section for more detailed information.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON SUGAR BEETS

- **DO NOT** apply more than 60 fl oz/A (*1.10 lbs ai/A*) of NULLIFY A/P per year.
- **DO NOT** apply NULLIFY A/P within 60 days of harvesting sugar beets.
- If a second application is needed, make the second application a minimum of 10 days after the first application.
- **DO NOT** exceed the single application rate maximum of 36 fl oz/A (*0.66 lb ai/A*).
- **DO NOT** make more than 2 applications per year.
- **DO NOT** plant rotation crops in a field treated with NULLIFY A/P within 120 days after the last application of this product with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale, which may be planted 70 days after the last application of this product. Corn, soybeans, canola, and sugar beets containing the LibertyLink trait may be planted at any time.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** apply NULLIFY A/P if sugar beets show injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.

APPLICATION DIRECTIONS FOR USE ON CANOLA

Apply NULLIFY A/P only to canola labeled as LibertyLink®. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

APPLICATION TIMING

Applications of NULLIFY A/P on canola may be made from the cotyledon stage up to the early bolting stage of the canola. Slight discoloration of the canola may be visible after application. This effect is temporary and will not influence crop growth, maturity, or yield.

Apply to small and actively growing weeds, targeting weeds less than 3 inches in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed control may be reduced when applications are made when heavy dew, fog and mist/rain are present or when weeds are under stress due to drought, cool temperatures or extended periods of cloudiness. NULLIFY A/P is a foliar-active material with little or no soil-residual activity. NULLIFY A/P is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment. For best result, on lambsquarters, Palmer amaranth and velvetleaf control, make applications of Glufosinate between dawn and 2 hours before sunset.

APPLICATION RATES

Apply NULLIFY A/P at 22 to 29 fl oz/A (0.40 - 0.53 lb ai/A) per application. If a second application of NULLIFY A/P is needed, make the second application in a minimum of 7 days after the first application. The maximum annual rate of NULLIFY A/P on canola is 87 fl oz/A (1.59. lbs ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations, use a minimum spray volume of 20 gallons per acre.

APPLICATION RATES WITH TANK MIX PARTNERS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Apply NULLIFY A/P at 22 to 29 fl oz (0.40 - 0.53 lb ai) per acre per application, depending on weed species, size and density per weed chart.

Tank mix partners advised to enhance grass control, including products containing quizalofop p-ethyl, sethoxydim and clethodim.

If a second application is needed, make the second application in a minimum of 7 days after the first application.

Tank mixes may aid in the performance of NULLIFY A/P. Please refer to weed chart tables for a listing of weed species controlled at this rate. No additional surfactant is needed with any tank mix partner. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Do not mix NULLIFY A/P mix with any product containing a label prohibition against such mixing.

ADJUVANTS

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 pounds per acre. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (such as temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds like lambsquarters and velvetleaf under difficult environmental conditions (such as low relative humidity) or hard water. The use of an anti-foam agent is advised.

SURFACTANTS/OILS

The use of additional surfactants or crop oils in tank mixes with NULLIFY A/P may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

NOZZLE SPRAY QUALITY

Use medium to coarse nozzles.

NULLIFY A/P is a contact herbicide and requires proper nozzles with uniform thorough spray coverage to achieve optimum weed control.

See **SPRAY DRIFT MANAGEMENT** section for more detailed information.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON CANOLA

1. **DO NOT** use on canola in the states of Alabama, Delaware, Georgia, Kentucky, Maryland, New Jersey, North Carolina, South Carolina, Tennessee, Virginia and West Virginia.
2. **DO NOT** apply more than two applications of NULLIFY A/P per year. Sequential applications must be at least 10 days apart.
3. **DO NOT** apply NULLIFY A/P within 65 days of harvesting canola.
4. **DO NOT** exceed the maximum single application rate of 43 fl oz/A (0.79 lb ai/A).
5. **DO NOT** apply more than 87 fl oz/A (1.59 lbs ai/A) of NULLIFY A/P per year.
6. **DO NOT** graze the treated crop or cut for hay.
7. **DO NOT** apply NULLIFY A/P if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
8. **DO NOT** apply this product through any type of irrigation system.
9. Refer to the "**Rotational Crop Restrictions**" section under the "**Product Information**" heading of this label for the appropriate rotational crop plant back intervals.

APPLICATION DIRECTIONS FOR CANOLA FOR LIBERTY LINK SEED PROPAGATION

Up to three applications of NULLIFY A/P at up to 29 fl oz/A (0.53 lb ai/A) per application may be made to canola for LibertyLink seed propagation. Applications may be made from the cotyledon stage up to the early bolting stage (e.g., BBCH 18-30, between just prior to stem elongation/bolting, eight or more leaves and beginning of stem elongation, no internodes).

RESTRICTIONS TO THE DIRECTIONS FOR CANOLA FOR LIBERTYLINK SEED PROPAGATION

- **DO NOT** apply more than three applications of NULLIFY A/P at up to 29 fl oz/A (0.53 lb ai/A) per application per year.
- Sequential applications must be made more than 10 days apart.
- **DO NOT** apply more than 87 fl oz/A (1.59 lbs ai/A) of NULLIFY A/P per year.
- **DO NOT** apply NULLIFY A/P beyond the early bolting stage or within 65 days of harvesting canola seed.
- **DO NOT** use treated canola seed for food, feed or oil purposes.
- **DO NOT** apply NULLIFY A/P if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.

APPLICATION DIRECTIONS FOR USE ON SWEET CORN*

*Not for use in CA

Apply NULLIFY A/P only to sweet corn containing the LibertyLink® trait. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

APPLICATION TIMING

Applications of NULLIFY A/P on sweet corn may be made from emergence until the V-6 stage of growth; i.e., 6 developed collars, whichever comes first.

Apply to small and actively growing weeds, targeting weeds less than 3 inches in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed control may be reduced when applications are made when heavy dew, fog and mist/rain are present or when weeds are under stress due to drought, cool temperatures or extended periods of cloudiness. NULLIFY A/P is a foliar-active material with little or no soil-residual activity. NULLIFY A/P is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment. For best results on lambsquarters, Palmer amaranth and velvetleaf, make applications of Glufosinate between dawn and 2 hours before sunset.

APPLICATION RATES

Apply at a rate of 22 fl oz/A (0.40 lb ai/A), depending on weed species, size and density per weed chart. If required, a second application of 22 fl oz/A (0.40 lb ai/A) can be applied. The second application must be made a minimum 7 days after the first application. The maximum annual rate of NULLIFY A/P on sweet corn is 44 fl oz/A (0.80 lb ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations, use a minimum spray volume of 20 gallons per acre.

APPLICATION RATES WITH TANK MIX PARTNERS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Apply NULLIFY A/P at 22 fl oz/A (0.40 lb ai/A) per application, depending on weed species, size and density per weed chart. Advised tank mix partners, including products containing atrazine, tembotrione, thiencazabonemethyl, and dicamba DGA salt.

If a second application is needed, make the second application in a minimum of 7 days after the first application.

Tank mixes may aid in the performance of NULLIFY A/P. Please refer to weed chart tables for a listing of weed species controlled at this rate.

No additional surfactant is needed with any tank mix partner. The tank mix partner must be used in accordance with the label limitations, restrictions and precautions. Do not exceed any labeled dosage rates. Do not mix NULLIFY A/P mix with any product containing a label prohibition against such mixing.

ADJUVANTS

Ammonium sulfate (AMS) may be used at 1.5 to 3.5 lb/A. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (such as temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds like lambsquarters and velvetleaf under difficult environmental conditions (such as low relative humidity) or hard water. The use of an anti-foam agent is advised.

SURFACTANTS/OILS

The use of additional surfactants or crop oils in tank mixes with NULLIFY A/P may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

NOZZLE SPRAY QUALITY

Use medium to coarse nozzles. NULLIFY A/P is a contact herbicide and requires proper nozzles with uniform thorough spray coverage to achieve optimum weed control.

See **SPRAY DRIFT MANAGEMENT** section for more detailed information.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON SWEET CORN

1. **DO NOT** apply NULLIFY A/P within 50 days of harvesting sweet corn ears and within 55 days of harvesting stover.
2. **DO NOT** apply more than 44 fl oz/A (0.80 lb ai/A) of NULLIFY A/P on sweet corn per year.
3. **DO NOT** apply more than two applications of NULLIFY A/P to sweet corn per year. Sequential applications must be at least 7 days apart.
4. **DO NOT** exceed the maximum single application rate of 22 fl oz/A (0.40 lb ai/A).
5. If NULLIFY A/P was used in a burndown application, no post emergence applications may be made to the crop.
6. **DO NOT** use nitrogen solutions as spray carriers. A silicone-based anti-foam agent may be added if needed.
7. **DO NOT** apply NULLIFY A/P if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall etc.).
8. **DO NOT** apply this product through any type of irrigation system.
9. Refer to the **Rotational Crop Restrictions** section under the **Product Information** heading of this label for the appropriate rotational crop plantback intervals.

See the **Application Methods for Broadcast Application, Mixing Instructions** and **Weed Control for Row Crops** tables on this label for further instruction.

APPLICATION DIRECTIONS FOR USE ON FIELD CORN AND SILAGE CORN

Apply NULLIFY A/P only to corn labeled as LibertyLink®. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

APPLICATION TIMING

Applications of NULLIFY A/P on corn may be made from emergence until the V-6 stage of growth, i.e., 6 developed collars, whichever comes first.

Apply to small and actively growing weeds, targeting weeds less than 3 inches in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed control may be reduced when applications are made when heavy dew, fog and mist/rain are present or when weeds are under stress due to drought, cool temperatures or extended periods of cloudiness. NULLIFY A/P is a foliar-active material with little or no soil-residual activity. NULLIFY A/P is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment. For best results on lambsquarters, Palmer amaranth and velvetleaf make applications of NULLIFY A/P between dawn and 2 hours before sunset

APPLICATION RATES

Apply NULLIFY A/P at 29 to 43 fl oz/A (0.53 - 0.79 lb ai/A) per application depending on weed species, size and density per weed chart. If a second application is needed, make the second application at up to 29 fl oz/A (0.53 lb ai/A) with a minimum of 7 days after the first application. The maximum rate of NULLIFY A/P on field corn and silage corn is 87 fl oz/A (1.59 lb ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as density canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

APPLICATION RATES WITH TANK MIX PARTNERS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Apply NULLIFY A/P at 29 to 43 fl oz/A (0.53 - 0.79 lb ai/A), depending on weed species, size and density per weed chart. Advised tank mix partners, including products containing atrazine, tembotrione, thien carbazonemethyl and dicamba, DGA salt. If a second application is needed, make the second application in a minimum of 7 days after the first application. Tank mixes may aid in the performance of NULLIFY A/P.

Please refer to weed chart tables for a listing of weed species controlled at this rate. No additional surfactant is needed with any tank mix partner. The tank mix partner must be used in accordance with the label limitations, restrictions and precautions.

Do not exceed any labeled dosage rates. Do not mix NULLIFY A/P mix with any product containing a label prohibition against such mixing.

ADJUVANTS

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 lb/A. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (such as temperature) and potential for leaf burn. AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (such as low relative humidity) or hard water. The use of an anti-foam agent is advised.

SURFACTANTS / OILS:

The use of additional surfactants or crop oils in tank mixes with NULLIFY A/P may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

NOZZLE SPRAY QUALITY

Use medium to coarse nozzles.

NULLIFY A/P is a contact herbicide and requires proper nozzles with uniform thorough spray coverage to achieve optimum weed control.

See **SPRAY DRIFT MANAGEMENT** section for more detailed information.

APPLICATION DROP NOZZLE EQUIPMENT

Applications of NULLIFY A/P Field corn and Corn Silage may be made with drop nozzles from emergence until corn is 36 inches tall. Avoid spraying into the whorl or leaf axils of the corn stalks

RESTRICTIONS TO THE DIRECTIONS FOR USE ON FIELD CORN AND SILAGE CORN

- **DO NOT** apply NULLIFY A/P within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder.
- **DO NOT** apply more than two applications per year. Sequential applications must be at least 10 days apart.
- **DO NOT** apply more than 87 fl oz/A (1.59 lb ai/A) of NULLIFY A/P on corn per year.
- **DO NOT** exceed the maximum single application rate of 43 fl oz/A (0.79 lb ai/A).
- **DO NOT** use nitrogen solutions as spray carriers. A silicone-based anti-foam agent may be added if needed.
- **DO NOT** apply NULLIFY A/P if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall etc.).
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the **“Rotational Crop Restrictions”** section under the **“Information”** heading of this label for the appropriate rotational crop plant back intervals.

APPLICATION DIRECTIONS FOR USE ON COTTON

Uniform, thorough spray coverage is necessary to achieve consistent weed control. NULLIFY A/P may be applied as a broadcast, over-the-top, post-emergence spray or as a directed spray only to LibertyLink® cotton. This product may be applied post-emergence to non-LibertyLink® cotton varieties or cultivars by using equipment designed to minimize contact of the spray with the cotton foliage. See the Application Methods on Non-LibertyLink® Cotton section for selection of shielding equipment. Severe injury or death may result if the NULLIFY A/P contacts the foliage or stems of cotton **NOT** labeled as LibertyLink®.

APPLICATION TIMING

Apply to small actively growing weeds, targeting weeds less than 3 inches in height. For additional information on weed heights, refer to the **WEED CONTROL FOR ROW CROPS** section.

Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed control may be reduced when applications are made when heavy dew, fog and mist/rain are present or when weeds are under stress due to drought, cool temperatures or extended periods of cloudiness. NULLIFY A/P is a foliar-active material with little or no soil-residual activity. NULLIFY A/P is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment. For best result, on lambsquarters, Palmer amaranth and velvetleaf control, make applications of Glufosinate between dawn and 2 hours before sunset.

APPLICATION RATES

Apply NULLIFY A/P to cotton from emergence up to the early bloom stage at 29 fl oz/A (0.53 lb ai/A) if environmental conditions prevent a timely herbicide application, a single application of up to 43 fl oz/A of NULLIFY A/P may be made to cotton.

If more than 29 fl oz/A (0.53 lb ai/A) are used in any single application, the yearly total may not exceed 72 fl oz/A (1.32 lb ai/A), including all application timings. See **Restrictions to the Directions for Use on Cotton** below for additional information.

Option 1: 3 post applications

Apply 29 fl oz/A (0.53 lb ai/A) per application depending on weed species, size and density per weed chart. If required a second application of 29 fl oz/A (0.53 lb ai/A) may be made 10-14 days after the first application. If required, a third application of 29 fl oz/A (0.53 lb ai/A) may be made 10-14 days after the second application. The yearly maximum rate of NULLIFY A/P on cotton is 87 fl oz/A (1.59 lb ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

Option 2: 2 post applications

Apply 32 to 43 fl oz/A (0.58 -0.79 lb ai/A) per application depending on weed species, size and density per weed chart. If required a second application of 29 fl oz/A (0.53 lb ai/A) can be applied. The sequential applications must be made minimum 10 days and may be made 14 days after each other. The maximum annual rate of NULLIFY A/P on cotton is 72 fl oz/A (1.32 lb ai/A) per acre.

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

Use Pattern	1st Application	2nd Application <i>Minimum 10 days up to 14 days after 1st application</i>	3rd Application <i>Minimum 10 days up to 14 days after 2nd application</i>	Yearly Maximum
Option 1	29 fl oz/A (0.53 lb ai/A)	29 fl oz/A (0.53 lb ai/A)	29 fl oz/A (0.53 lb ai/A)	87 fl oz/A (1.59 lbs ai/A)
Option 2	32 - 43 fl oz/ A (0.58-0.79 lb ai/A)	29 fl oz/A (0.53 lb ai/A)	None	72 fl oz/A (1.32 lbs ai/A)

TANK MIX ON COTTON

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Certain herbicide tank mixes may aid in the performance of NULLIFY A/P. NULLIFY A/P may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the cotton to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. NULLIFY A/P cannot be mixed with any product containing a label prohibition against such mixing.

ADJUVANTS

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 lb/A. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (such as temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds like lambsquarters and velvetleaf under difficult environmental conditions (such as low relative humidity) or hard water. The use of an anti-foam agent is advised.

SURFACTANTS / OILS

The use of additional surfactants or crop oils in tank mixes with NULLIFY A/P may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

NOZZLE SPRAY QUALITY

Use medium to coarse nozzles. NULLIFY A/P is a contact herbicide and requires proper nozzles with uniform thorough spray coverage to achieve optimum weed control.

See **SPRAY DRIFT MANAGEMENT** section for more detailed information.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON COTTON

- **DO NOT** apply NULLIFY A/P to cotton in Florida, South of Tampa (Florida Route 60), or in Hawaii, except for test plots or breeding nurseries.
- **DO NOT** apply NULLIFY A/P within 70 days prior to cotton harvest.
- Up to 3 applications of NULLIFY A/P may be made to cotton per year at a maximum application rate of 29 fl oz/A (0.53 lb ai/A).
- **DO NOT** apply more than 87 fl oz (1.59 lb ai/A) (including all application timings) to cotton per year under this application scenario. Sequential applications must be at least 10 days apart.
- If environmental conditions prevent timely applications resulting in large weeds or heavy infestations, a single application of NULLIFY A/P at up to 43 fl oz/A (0.79 lb ai/A) may be made to cotton.
- **DO NOT** apply more than 43 fl oz/A (0.79 lb ai/A) of NULLIFY A/P in a single application under this use scenario. If a single application greater than 29 fl oz (0.53 lb ai) is made, a subsequent application not to exceed 29 fl oz (0.53 lb ai) may be made to cotton. The annual total use rate under this scenario may not exceed 72 fl oz/A (1.32 lb ai) of NULLIFY A/P. Sequential applications must be at least 10 days apart.
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the **"Rotational Crop Restrictions"** section under the **"Product Information"** heading of this label for the appropriate rotational crop plant back intervals.

APPLICATION METHODS FOR NON-LIBERTYLINK COTTON

Application of NULLIFY A/P to cotton varieties not labeled as LibertyLink® requires the use of hooded spray equipment designed to minimize exposure of the spray to the cotton stand. A hooded sprayer directs the spray onto weeds, while shielding the cotton stand from contact. Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Extreme care must be exercised to avoid exposure of the desirable vegetation to the spray.

With a hooded sprayer, the spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground. If the hoods are raised, spray particles may escape and come into contact with the cotton, causing damage or destruction of the crop.

Herbicide rates and spray volume instructions are presented as broadcast equivalents and must be reduced in proportion to the area actually treated. Use the following formulas to calculate the correct rate and volume per planted (field) acre:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per acre} = \text{Amount of banded product needed per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast spray VOLUME per acre} = \text{Banded Spray Volume needed per acre}$$

TANK MIX ON COTTON

Certain tank mixes may aid in the performance of NULLIFY A/P. No additional surfactant is needed with any tank mix partner. NULLIFY A/P may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the cotton to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. NULLIFY A/P cannot be mixed with any product containing a label prohibition against such mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

POST-HARVEST / FALL BURNDOWN ON COTTON

NULLIFY A/P may be applied as a post-harvest burndown treatment to fields (after cotton harvest). Up to 43 fl oz/A (0.79 lb ai/A) of NULLIFY A/P may be applied in a single application to control larger weeds growing in the crop at the time of harvest. If more than 29 fl oz/A (0.53 lb ai/A) is used in a single application, the yearly total may not exceed 72 fl oz/A (1.32 lb ai/A), including all application timings. Refer to the **Rotational Crop Restrictions** section of this label for appropriate rotational crop information.

TANK MIX ON COTTON

Certain tank mixes may aid in the performance of NULLIFY A/P. No additional surfactant is needed with any tank mix partner. NULLIFY A/P may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the cotton to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. NULLIFY A/P cannot be mixed with any product containing a label prohibition against such mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

RESTRICTIONS TO THE DIRECTIONS FOR COTTON POST HARVEST BURNDOWN USE

- **DO NOT** apply more than 43 fl oz/A (0.79 lb ai/A) of NULLIFY A/P in a single application
- **DO NOT** apply more than 72 fl oz/A (1.32 lb ai/A) per year, including all application timings.
- **DO NOT** make more than two applications per year. **DO NOT** make second application within 10 days of first.

APPLICATION DIRECTIONS FOR USE ON SOYBEANS

Apply NULLIFY A/P only to soybeans designated as LibertyLink®. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

APPLICATION TIMING

Applications of NULLIFY A/P on soybeans may be made from emergence up to but not including the R1 bloom growth stage.

Apply to small actively growing weeds, targeting weeds less than 3 inches in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed control may be reduced when applications are made when heavy dew, fog and mist/rain are present or when weeds are under stress due to drought, cool temperatures or extended periods of cloudiness. NULLIFY A/P is a foliar-active material with little or no soil-residual activity. NULLIFY A/P is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment. For best results on lambsquarters, Palmer amaranth and velvetleaf, make applications of Glufosinate between dawn and 2 hours before sunset.

APPLICATION RATES

Apply NULLIFY A/P at 29 to 43 fl oz/A (0.53 - 0.79lb ai/A) depending on weed species, size and density per weed chart.

If a second application is needed, make the second application of 29 to 43 fl oz/A (0.53 - 0.79 lb ai/A), can be applied up to a yearly maximum of 87 fl oz/A (1.59 lbs ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (such as dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

Use Pattern Rate Ranges

1st Application	2nd Application Minimum of 5 days after 1st Application	Yearly Maximum
29-48 fl oz/A (0.53 - 0.79 lb ai/A)	29-43 fl oz/A (0.53-0.79 lb ai/A)	87 fl oz/A (1.59 lbs ai/A)

SOYBEAN TANK MIX INSTRUCTIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Certain herbicide tank mixes may complement NULLIFY A/P. No additional surfactant is needed with any tank mix partner. NULLIFY A/P may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the soybean to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. NULLIFY A/P cannot be mixed with any product containing a label prohibition against such mixing.

ADJUVANTS

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 lb/A. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (such as temperature) and potential for leaf burn. AMS has shown to improve weed control of difficult-to-control weeds like lambsquarters and velvetleaf under difficult environmental conditions (such as low relative humidity) or hard water. The use of an anti-foam agent is advised.

SURFACTANTS / OILS

The use of additional surfactants or crop oils in tank mixes with NULLIFY A/P may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

NOZZLE SPRAY QUALITY

Use medium to coarse nozzles. NULLIFY A/P is a contact herbicide and requires proper nozzles with uniform thorough spray coverage to achieve optimum weed control.

See **SPRAY DRIFT MANAGEMENT** section for more detailed information.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON SOYBEANS

- **DO NOT** apply NULLIFY A/P within 70 days of harvesting soybean seed.
- **DO NOT** apply more than 87 fl oz/A (1.59 lb ai/A). of NULLIFY A/P on soybeans per year.
- **DO NOT** apply more than 43 fl oz/A (0.79 lb ai/A). of NULLIFY A/P in a single application.
- **DO NOT** make more than 3 applications per year.
- **DO NOT** make sequential applications within 10 days of previous application.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** use nitrogen solutions as spray carriers. A silicone-based anti-foam agent may be added if needed.
- **DO NOT** apply NULLIFY A/P if soybeans show injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the **"Rotational Crop Restrictions"** section under the **"Product Information"** heading of this label for the appropriate rotational crop plant back intervals.
- Sequential applications must be at least 5 days apart

APPLICATION DIRECTIONS FOR CANOLA, CORN, COTTON, AND SOYBEAN SEED PROPAGATION

NULLIFY A/P may be applied to select out susceptible "segregates," i.e., canola, corn, cotton, and soybean plants that are sensitive to glufosinate-ammonium (i.e. do not contain the LibertyLink trait) during seed propagation.

- **Canola:** NULLIFY A/P may also be used in canola seed propagation as a foliar spray to selectively eliminate canola plants that do not carry the LibertyLink trait and as such, can be applied to remove susceptible segregates during canola seed propagation. Breeding material not possessing the LibertyLink trait will be severely injured or killed if treated with this herbicide.

See **Application Use Directions for Use on Canola** for use rates, application timing and use restrictions.

- **Corn:** Inbred lines, plants not possessing the LibertyLink trait, will be severely injured or killed if treated with this herbicide. A hooded sprayer may be used to protect plants from coming into contact with the herbicide application. For the selection of non-sensitive corn "segregates", NULLIFY A/P may be applied at 22 fl oz/A (0.40 lb ai/A) plus AMS at 3 lb/A (17 lb/100 gallons) when corn is in the V-3 to V-4 stage of growth, i.e., 3 to 4 developed collars. A second treatment of 22 fl oz/A (0.40 lb ai/A) plus AMS at 3 lb/A may be applied when the corn is in the V-6 to V-7 stage of growth or up to 24" tall. Sequential applications need to be at least 10 days apart. When temperatures exceed 85°F, the rate of AMS can be reduced to 1.5 lbs/A (8.5 lb/100 gallons) to reduce potential leaf burn.

See **Application Use Directions for Use on Corn** for use rates, application timing and use restrictions.

- **Cotton:** NULLIFY A/P may also be used in cotton seed propagation as a foliar spray to selectively eliminate cotton plants that do not carry the LibertyLink trait and as such, can be applied to remove susceptible segregates during cotton seed propagation. Breeding material not possessing the LibertyLink trait will be severely injured or killed if treated with this herbicide.

See **Application Use Directions for Use on Cotton** for use rates, application timing and use restrictions.

- **Soybean:** For the selection of non-sensitive soybean "segregates", NULLIFY A/P may be applied at up to 22 - 36 fl oz/A (0.40 - 0.66 lb ai/A) when soybean is in the third trifoliolate stage. A second treatment of 22 - 29 fl oz/A (0.40 - 0.53 lb ai/A) may be applied up to but not including the bloom growth stage of soybean. Sequential applications must be at least 5 days apart.

See **Application Use Directions for Use on Soybeans** for use rates, application timing and use restrictions.

APPLICATION DIRECTIONS FOR USE ON LISTED TREE, VINE, AND BERRY CROPS

Apply NULLIFY A/P to the tree, vine, and berry crops listed below. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

REGISTERED CROPS

BERRIES:

Crop Subgroup 13-B: Bushberry subgroup

Blueberry, highbush; blueberry, lowbush; currant; elderberry; gooseberry; huckleberry

Juneberry; lingonberry; salal

CITRUS CROP GROUP 10-10:

Orange or tangerine/mandarin, Calamondin; citron, citrus hybrids; Mediterranean Mandarin; orange, sour; orange, sweet; satsuma darin; tachibana orange; tangerine (mandarin); tangelo; tangor, trifoliate orange; cultivars, varieties and/or hybrids of these

Lemon or lime - Australian desert lime; Australian finger lime; Australian round lime; brown river finger lime; kumquat; lemon; lime; mount white lime; New Guinea wild lime; Russel River lime; sweet lime; Tahiti lime; cultivars, varieties and/or hybrids of these

Grapefruit - Grapefruit; Japanese summer grapefruit; pummelo; tangelo; uniq fruit; cultivars, varieties and/or hybrids of these.

OLIVES: All olive varieties

POME FRUIT (CROP GROUP 11-10):

Crop Group 11. Pome Fruits Group

Apple; crabapple; loquat; mayhaw; pear; pear, oriental; quince; azarole; hook; medlar; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties and/or hybrids of these

STONE FRUIT (CROP GROUP 12-12):

Crop Group 12. Stone Fruits Group

Apricot; cherry, sweet; cherry, tart; nectarine, peach; plum; plum, chickasaw; damson; plum, Japanese; plumcot; prune; capulin; jujube and sloe; cultivars, varieties and/or hybrids of these.

TREE NUTS (CROP GROUP 14 INCLUDING PISTACHIOS):

Crop Group 14. Tree Nuts Group

Almond; beech nut; Brazil nut; butternut; cashew; chestnut; chinquapin; filbert (hazelnut), hickory nut, macadamia nut (bush nut), pecan, pistachios, and walnut, black and English

GRAPES: all grape varieties (table, wine, and raisins)

APPLICATION TIMING

For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of NULLIFY A/P. Weed Control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. Do not retreat these weeds with NULLIFY A/P until sufficient regrowth has occurred.

Apply NULLIFY A/P as a directed spray to control undesirable vegetation in tree, vine and berries listed on this label. Apply as a broadcast, banded, or spot treatment application depending on the situation to control weeds listed under the heading "Weeds Controlled in Tree, Vine and Berry crops". Avoid direct spray or drift to desirable vegetation. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of NULLIFY A/P may be necessary to control plants generating from underground parts or seed.

Avoid contact of NULLIFY A/P solution, spray, drift or mist with green bark, stems, or foliage, as injury may occur to trees and vines. **Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes or waxed containers. Contact of NULLIFY A/P with parts of trees or vines other than mature brown bark can result in serious damage.**

APPLICATION METHODS FOR BROADCAST APPLICATIONS

Apply NULLIFY A/P at the rates listed below for broadcast applications based on weed size and stage of growth.

Weed Size and Stage	NULLIFY A/P Rate	
	fl oz/A	lb ai/A
Weeds < 3" in height	48 fl oz/A	0.88 lb ai/A
Weeds < 6" in height pre-tiller grasses	56 fl oz/A	1.02 lb ai/A
Weeds > 6" in height and/or grasses that have tillered	56 - 82 fl oz/A	1.02 - 1.50 lbs ai/A

APPLICATION METHODS FOR BANDED SPRAY APPLICATIONS

Banded applications may be used using the following formula to calculate the amount of herbicide needed for orchard or vineyard strip sprays:

Band width in inches	X Rate per acre	= Amount of herbicide
Row width in inches	broadcast	needed for treatment

APPLICATION METHODS FOR SPOT OR DIRECTED-SPRAY APPLICATIONS

For spot or directed spray application, mix NULLIFY A/P at 1.7 fl oz (0.03 lb ai) of product per gallon of water. Apply to undesirable vegetation foliage until wet but prior to runoff. Ensure uniform and complete coverage. Thoroughly clean the sprayer following use. DO NOT make spot or directed spray applications to tree or vine trunk as injury may occur.

WEEDS CONTROLLED IN TREE, VINE AND BERRY CROPS

Broadleaf Weeds

Alkali sida	Goosefoot	Morningglory, ivyleaf	Shepherd's-Purse
Ammannia, purple	Gromwell, field	Morningglory, pitted	Smartweed, Pennsylvania
Arrowhead, California	Groundcherry, cutleaf	Mullein, turkey	Sowthistle, annual
Buckwheat, wild	Groundsel, common	Mustard, wild	Spurge, prostrate
Buffalobur	Henbit	Nettle	Starthistle, yellow
Burclover, California	Jimsonweed	Nightshade, black	Sunflower, common
Carpetweed	Knotweed	Nightshade, eastern black	Sunflower, prairie
Chickweed, common	Kochia	Nightshade, hairy	Sunflower, volunteer
Chinese thornapple	Lambsquarters, common	Pennycress	Swinecress
Cocklebur, common	Lettuce, miner's	Pigweed, redroot	Thistle, Russian
Copperleaf, Virginia	Lettuce, prickly	Pineapple-weed	Turnip, wild
Cudweed	London rocket	Puncturevine	Velvetleaf
Cutleaf eveningprimrose	Mallow, common	Purslane, common	Vervain
Dodder	Malva (little mallow)	Radish, wild	Vetch
Eclipta	Marestail	Ragweed, common	Virginia copperleaf
Fiddleneck	Mayweed	Ragweed, giant	Willowherb, panicle
Filaree	Morningglory,	Redmaids	
Filaree, redstem	entireleaf		
Fleabane, annual			

Grass Weeds

Barnyardgrass	Crabgrass, smooth	Junglerice	Shattercane
Bluegrass, annual	Cupgrass, woolly	Oat, wild	Sprangletop
Brome, riggut	Foxtail, giant	Panicum, fall	Stinkgrass
Bromegrass, downy	Foxtail, green	Panicum, Texas	Wheat, volunteer
Canarygrass	Foxtail, yellow	Rush, toad**	Windgrass
Chess, soft	Goosegrass	Ryegrass, annual*	Witchgrass
Crabgrass, large	Johnsongrass, seedling	Sandbur, field	

Biennial and Perennial Weeds

Aster, white heath	Dallisgrass	Mullein, common	Rocket, yellow
Bindweed, field	Dandelion	Mustard, tansy	Rose, wild
Bindweed, hedge	Dock, curly	Nutsedge, purple	Rubus spp.
Bluegrass, Kentucky	Dogbank (hemp)	Nutsedge, yellow	Spurge, leafy
Bromegrass, smooth	Fescue	Onion, wild	Thistle, bull
Bulrush**	Goldenrod, gray	Orchardgrass	Thistle, musk
Burdock	Guineagrass	Paragrass	Torpedograss
Canada thistle	Horsetail	Plantain	Vaseygrass
Clover, Alsike	Lovegrass	Poison ivy/oak	Woodsorrel
Clover, red	Mugwort	Quackgrass	Yarrow, common
Clover, white			

* apply to annual ryegrass prior to 3 inches in height

** indicates suppression

RESTRICTIONS TO THE DIRECTIONS FOR USE ON TREE, VINE, AND BERRY CROPS

- **DO NOT** apply more than 164 fl oz of NULLIFY A/P per acre (3.0 lbs ai/A) to berry bushes and stone fruit in a 12-month period.
- **DO NOT** exceed the maximum single application rate of 82 fl oz/A (1.50 lb ai/A)
- **DO NOT** make more than 2 applications per year at a maximum rate of 82 fl oz/A (1.50 lb ai/A) per application to berry bushes and stone fruit.
- **DO NOT** apply more than 246 fl oz (4.50 lbs ai/A) of this product per acre to tree nuts, vines, pome fruits, citrus and olives in any calendar year.
- **DO NOT** make more than 3 applications at a maximum rate of 82 fl oz/A (1.50 lb ai/A) per application to tree nuts, vines, pome fruits, citrus and olives.
- **DO NOT** graze, harvest, and/or feed treated orchard cover crops to livestock.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product aerially to tree, berry, or vine crops.
- **DO NOT** apply this product within 14 days of nut, fruit, berry or grape harvest.
- Applications to citrus fruits, pome fruits and olives must be a minimum of 14 days apart.
- Applications to stone fruit must be a minimum of 28 days apart.
- Applications to berry bushes must be a minimum of 14 days apart.
- **DO NOT** make spot spray applications to suckers, as tree injury may occur.

SUCKER CONTROL WITH NULLIFY A/P HERBICIDE

NULLIFY A/P will reduce or eliminate sucker growth when applied to suckers that are young, green and uncaloused. For sucker control, apply a split application approximately 4 weeks apart at 56 fl oz/A (1.02 lb ai/A). Coverage of all sucker foliage is necessary for optimum control. Suckers should not exceed 12 inches in length.

TANK MIX PARTNER

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

NULLIFY A/P does not provide residual weed control or control of unexposed plant parts. Certain herbicide tank mixes may aid in the performance of NULLIFY A/P or be added to provide residual herbicide activity. No additional surfactant is needed with any tank mix partner. NULLIFY A/P may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. NULLIFY A/P cannot be mixed with any product containing a label prohibition against such mixing.

diuron	napropamide	oryzalin	terbacil
flumioxazin	norfluzon	simazine	

APPLICATION DIRECTIONS FOR POTATO VINE DESICCATION

APPLICATION RATES AND TIMING

Apply NULLIFY A/P at the beginning of natural senescence of potato vines. Apply 21 fl oz/A (0.38 lb ai/A). Do not split this application or apply more than one application per harvest. Potato varieties with heavy or dense vines may require an application of another desiccation product to complete vine desiccation.

Thorough coverage of the potato vines to be desiccated is essential. Use a sufficient volume of water (20 to 100 gpa) to obtain a thorough coverage of the potato vines. Vary the gallons of water per acre and the spray pressure as indicated by the density of the potato vines to assure thorough spray coverage. Increase the spray volume to at least 30 gallons of water per acre when the potato vine canopy is dense or under cool and dry conditions. Apply NULLIFY A/P with the spray boom as low as possible to achieve thorough coverage of the potato vines for best control and to minimize drift potential.

RESTRICTIONS TO THE DIRECTIONS FOR USE IN POTATO VINE DESICCATION

- **DO NOT** apply more than 21 fl oz/A (0.38 lb ai/A) to potato vines per year or per single application.
- **DO NOT** harvest potatoes until 9 days or more after application of NULLIFY A/P.
- **DO NOT** apply to potatoes grown for seed.
- Potatoes, canola, corn, cotton, soybean, and sugar beets may be planted at any time after the application of NULLIFY A/P as a potato vine desiccant.
- **DO NOT** plant treated areas to wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale until 30 or more days after an application of NULLIFY A/P as a potato vine desiccant.
- **DO NOT** plant treated areas to crops other than those listed in this use precautions section until 120 or more days after an application of NULLIFY A/P as a potato vine desiccant.
- **DO NOT** split this application or apply more than one application per harvest.

FALLOW FIELDS OR POST HARVEST

NULLIFY A/P may be used as a substitute for tillage in fallow fields to control or suppress weeds listed in the **Weed Control for Row Crops** section of this label. Applications may be made in fallow fields, post-harvest, prior to planting or emergence of any crop listed on this label. Apply NULLIFY A/P at 22 or 29 fl oz/A (0.40 - 0.53 lb ai/A) to fallow fields to control specific weeds. NULLIFY A/P must be applied with ammonium sulfate. Tank mixes with 2,4-D, glyphosate or atrazine are advised with NULLIFY A/P to enhance total weed control. When using NULLIFY A/P in tank mix combinations, follow the precautions and directions of use of the most restrictive label. See the **Application and Mixing Procedures** section of this label for additional information on how to apply this product. See the **"Product Information"** section of this label for rotational crop restrictions.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON FALLOW FIELDS OR POST HARVEST

- **DO NOT** apply more than 29 fl oz/A (0.53 lbs ai/A) in a single application.
- **DO NOT** make more than 3 applications per year
- **DO NOT** make sequential applications sooner than 14 days apart.
- **DO NOT** apply more than 87 fl oz/A (1.59 lbs ai/A) per year.

NON-CROP USES

NULLIFY A/P controls annual and perennial weeds in non-crop areas defined below in the "Where to Apply Section". Applications may be made on a broadcast, banded or spot treatment basis depending on the situation. Avoid direct spray or drift to desirable vegetation. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat treatments may be necessary to control plants generating from underground parts or seed.

WHEN TO APPLY

NULLIFY A/P is a foliar-active material. Best results are obtained when weeds are actively growing. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application of the highest rate directed.

NULLIFY A/P must be applied at the labeled rate in the **HOW TO APPLY** section. Repeat applications of NULLIFY A/P or tank mixes of NULLIFY A/P plus one or more appropriate residual herbicide(s) listed on this label will be needed to control weeds emerging from underground parts or seeds.

HOW TO MIX

NULLIFY A/P must be mixed with water to make finished spray solution as follows:

1. Fill the spray tank with the required amount of water.
2. Add the proper amount of product, then mix thoroughly.

HOW TO APPLY

Spot or Directed Applications

This product may be used as a spot or directed spray application using 0.4 to 0.75 fl oz/gal of water (0.007 - 0.014 lbs ai/gal of water) of water depending upon the weed and stage of growth as shown in the following sections. Spray undesirable vegetation foliage on a spray-to-wet basis. Do not apply beyond runoff. Ensure uniform and complete coverage. Use a coarse spray. Do not spray during windy conditions. Backpack, pump-up, and hydraulic sprayers may be used. Thoroughly clean the sprayer following use.

When making spot treatments **DO NOT** exceed broadcast per acre use rates.

Broadcast or Boom Applications

Apply 12 - 38 fl oz/A (0.22 - 0.69 lb ai/A) depending upon the weed and stage of growth as shown in the following sections. Use a minimum of 40 gallons of water per acre with a minimum of 30-psi spray pressure.

Aerial Applications

Apply as a foliar treatment using a minimum of 5 gallons of water per acre to ensure thorough coverage. Do not apply when winds are gusty or under conditions which favor drift on to desirable vegetation. Applications under conditions which cause drift of this product will result in damage to any vegetation contacted. Drift control additives may be used. If a drift control additive is used, observe and follow all directions and precautions as specified on the additive label.

Tank Mix Directions for Non-crop Uses

NULLIFY A/P is compatible in tank mixes with many other herbicides including non-selective herbicides including glyphosate.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank mix applications of NULLIFY A/P plus the following herbicides are advised for broad-spectrum post emergence and preemergence weed control:

Isopropylamine salt of imazapyr	butroxydim	norflurazon
prodiamine	isoxaben	Diglycolamine salt of 3,6-dichloro-o-anisic Acid
oryzalin	pendimethalin	oxadiazon

A compatibility test must be conducted with any potential tank mix partner with NULLIFY A/P, except with any one of those listed above. Using a clear glass quart jar, conduct the test as described below:

1. Fill the jar three-quarters full with water.
2. Add the appropriate amount of herbicide in the following order: (a) dry flowable, (b) wettable powder, (c) aqueous suspensions, (d) flowables, (e) liquids and (f) solutions and emulsifiable or liquid concentrates. Shake or gently stir jar after each addition to thoroughly mix.
3. After adding all ingredients, let the mixture stand for 15 minutes and then look for separation, large flakes, precipitates, gels, and heavy oily film on the jar or other signs of incompatibility.
4. If the compatibility test shows signs of incompatibility, do not tank mix the product tested with NULLIFY A/P.

For the Following Weeds Controlled by NULLIFY A/P Apply:

Spot application:

Apply 0.75 fl oz/gal of water (*0.014 lb ai/gal of water*) when the weed height or diameter is less than 6 inches.
Apply 1.25 fl oz/ gal of water (*0.023 lb ai/gal of water*) when the weed height or diameter is 6 inches or greater.

Broadcast application:

Apply 40 fl oz/A (*0.73 lb ai/A*) when the weed height or diameter is less than 6 inches.
Apply 56 fl oz/A (*1.02 lb ai/A*) when the weed height or diameter is 6 inches or greater.

Broadleaf Weeds

Chickweed	Jimsonweed	Marestail
Clover	Kochia	Purslane
Common	London rocket	Shepherdspurse
Cocklebur	Malva(little mallow)	Smartweed
Filaree		

Grasses and Sedges

Barnyardgrass	Cupgrass	Green Foxtail	Smallflower Alexandergrass
Fall Panicum		Johnsongrass (rhizome)	Lovegrass (Signalgrass)
Giant		Shattercane	Stinkgrass
Foxtail			Windgrass, yellow
Goosegrass			Foxtail

For the Following Weeds Controlled by NULLIFY A/P Apply:

Spot application:

Apply 1.25 fl oz/gal of water (*0.023 lb ai/gal of water*) when the weed height or diameter is less than 6 inches.

Apply 1.75 fl oz/gal of water (*0.032 lb ai/gal of water*) when the weed height or diameter is 6 inches or greater.

Broadcast application:

Apply 56 fl oz/A (*1.02 lbs ai/A*) when the weed height or diameter is less than inches tall. Apply 80 fl oz/A (*1.46 lbs ai/A*) when the weed height or diameter is 8 inches or greater.

Broadleaf weeds

Annual sowthistle	Lambsquarter	Tansy mustard
Bindweed	Leafy spurge	Velvetleaf
Buffalorburr	Mugwort	Vervain
Burdock	Musk thistle	Virginia copperleaf
Canada thistle	Nettle	White heath aster
Curly dock	Nightshade	Wild buckwheat
Dandelion	Pennycress	Wild mustard
Dogbane (hemp)	Pigweed, redroot	Wild onion
Field growwell	Plantain	Wild rose
Fleabane	Prickly lettuce	Wild turnip
Goldenrod	Ragweed	Wood sorrel
Horsetail	Russian thistle	Yellow rocket

Grasses and Sedges

Annual bluegrass	Downy bromegrass	Ryegrass
Bahiagrass	Fescue	Sandbur
Barley	Guineagrass	Smooth bromegrass
Bermudagrass	Kentucky bluegrass	Torpedograss
Carpetgrass	Nutsedge	Vaseygrass
Crabgrass	Paragrass	Wheat
Dallisgrass	Quackgrass	Wild oat

Additional Use Directions

1. Use higher rates within the directed rate range for plant sizes listed when vegetation cover is dense or when weeds are growing under stressed conditions such as drought or when average temperatures are below 50°F.
2. The addition of 8.5 to 17 pounds of ammonium sulfate (spray grade) per 100 gallons of water (1 to 2% by weight) or 2 to 4 pounds of ammonium sulfate per acre may improve the level of weed control.

Use on Woody Species (Not For Use in California)

When applied as labeled, NULLIFY A/P will provide control, partial control, or suppression of certain perennial woody weed species. Apply 64 -192 fl oz/A (1.19 - 3.51 lb ai/A). Use the higher specified rates per acre of this product when conditions are not optimum for spray penetration, such as when vegetation growth is heavy or dense. Lower specified rates may be used when the target species is a conifer and when vegetation growth conditions allow for uniform spray coverage.

Blackberry	<i>Rubus</i> spp.
Deer brush	<i>Ceanothus integerrimus</i>
Douglas fir	<i>Pseudotsuga menziesii</i>
Gallberry	<i>Ilex</i> spp.
Hazel	<i>Corylus</i> spp.
Honeysuckle	<i>Lonicera</i> spp.
Huckleberry	<i>Gaylussacia</i> spp.
Maple	<i>Acer</i> spp.
Multiflora rose	<i>Rosa multiflora</i>
Oak	<i>Quercus</i> spp.
Pine	<i>Pinus</i> spp.
Poison ivy	<i>Toxicodendron radicans</i>
Poison oak	<i>Toxicodendron toxicarium</i>
Roundleaf greenbrier	<i>Smilax rotundifolia</i>
Salmonberry	<i>Rubus spectabilis</i>
Sweet gum	<i>Liquidambar styraciflua</i>
Sumac	<i>Rhus</i> spp.
Thimbleberry	<i>Rubus parviflorus</i>
Trumpetcreeper	<i>Campsis radicans</i>
Vine maple	<i>Acer circinatum</i>
Western red cedar	<i>Thuja plicata</i>

WHERE TO APPLY

Trimming and Edging

NULLIFY A/P may be used for trimming and edging landscape areas including: around individual trees and shrubs, landscape beds, foundations, fences, driveways, paths, and parking areas; also on golf courses along cart paths, around sign and light posts, and around sand traps. For control of weeds emerging from seed, the use of NULLIFY A/P in a tank mix with preemergence herbicides is advised. If spraying in areas adjacent to desirable plants, use a shield made of cardboard, plywood, or sheet metal while spraying to help prevent spray from contacting foliage of desirable plants. Refer to the How to Apply section of this labeling for appropriate application rates to control specific weeds.

Farmsteads, Recreational and Public Areas

When applied as a spot or directed spray application, this product controls annual and perennial weeds listed on this label in areas including: areas around farmstead building foundations, shelter belts, along fences, airports, commercial plants, storage and lumber yards, educational facilities, fence lines, ditch banks, dry ditches, roadsides, schools, parking lots, tank farms, pumping stations, and parks. Refer to the How to Apply section of this labeling for appropriate application rates to control specific weeds.

Dormant Bermudagrass (Not for use on Residential Turf/Turfgrass/Lawns)

NULLIFY A/P may be used to control winter annual weeds in well-established ornamental dormant hybrid or common Bermudagrass. Apply only when the turf is fully dormant and prior to spring green-up or severe turfgrass injury or delayed green-up may occur. For best results, apply NULLIFY A/P at a rate of 40 - 80 fl oz/A (0.73 - 1.46 lb ai/A) after most weeds have germinated and are in an early growth stage. Refer to the Weeds Controlled by NULLIFY A/P section of this label for selecting specified rates. Applications of NULLIFY A/P may also be used to suppress or control undesirable biennial or perennial weeds. Do not apply more than 80 fl oz (1.46 lbs ai) of NULLIFY A/P per acre per year for this use. Avoid high volume and spot applications where spray volume exceeds 80 gallons per acre or injury or delayed greenup may occur.

Ornamentals and Christmas Trees

When applied as specified by this label, this product may be used for the control of undesirable vegetation in site preparation prior to planting, around and within shade and greenhouses, and as a directed spray around containers and field-grown established ornamentals and Christmas trees.

DO NOT apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation or injury may result.

DO NOT apply NULLIFY A/P as an over-the-top broadcast spray in ornamentals and shade or Christmas trees.

Directed spray application:

NULLIFY A/P may be applied as a directed spray to control in-row weeds in field-grown woody plants. Refer to the How to Apply section of this labeling for appropriate application rate to control specific weeds. This product may also be used between and around containers and in site preparation for new planting.

Site preparation application:

This product may be used for pre-plant site preparation for the control of annual and perennial weeds listed on this label, in ornamental and Christmas tree plantings. Ornamentals and Christmas trees may be planted into the treated area after the restricted entry interval (REI) of 12 hours has elapsed. Refer to the How to Apply section of this labeling for appropriate application rates to control specific weeds.

Greenhouse and shade house applications:

NULLIFY A/P may be used to control weeds in greenhouses and shade- houses. Air circulation fans must be turned off during application. Apply NULLIFY A/P as a directed spray, using large droplet and low-pressure type nozzles. Avoid drift and direct contact with desirable vegetation. Do not use in greenhouses or shade houses containing edible crops.

USE RESTRICTIONS FOR NON-CROP USE

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with this product.
- **DO NOT** exceed maximum use rate of 80 fl oz/A (*1.46 lb ai/A*) per single application for broadcast or boom applications.
- **DO NOT** make more than 3 applications per year for broadcast or boom applications but no more than 2 applications per year on Dormant bermudagrass.
- **DO NOT** exceed maximum use rate of 1.75 fl oz/gal of water (*0.032 lbs ai/gal of water*) for spot or directed applications and do not apply beyond runoff.
- **DO NOT** apply more than 240 fl oz (*4.50 lbs ai/A*) of this product per acre per year to non-crop areas except on Dormant Bermudagrass do not apply more than 80 fl oz per acre per year.
- Applications must be made at least 14 days apart in non-crop areas.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not use or store near heat or open flame. Keep the container tightly closed and dry in a cool, well-ventilated place. Storage temperature must not exceed 125° F. If storage temperature for bulk NULLIFY A/P is below 32° F, the material must not be pumped until its temperature exceeds 32° F. Protect against direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER HANDLING:

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once container is rinsed, then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill, or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

All refillable container types (containers with capacities greater than 50 lbs)

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. This is a sealed returnable container to be used only for NULLIFY A/P. When this container is empty, it must not be opened, cleaned, or discarded. Empty containers must be returned to the original purchase location.

Bottom discharge Intermediate Bulk Container (IBC) (containers with capacities greater than 50 lbs)

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Empty the remaining contents from the Intermediate Bulk Container (IBC) into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inch on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve. Contact your Ag retailer or Control Solutions, Inc. for container return, disposal and recycling directions.

SEED DISPOSAL: To dispose of out-of-date or otherwise unmarketable seed from plants which have been treated with NULLIFY A/P, broadcast and lightly incorporate seed into field soils using disc or other suitable implement. Any resulting crop may be destroyed by chemical or mechanical means. Alternatively, seed may be destroyed by deep burial, incineration or landfill disposal.

IMPORTANT READ BEFORE USE

Read the entire Directions for Use Conditions Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable return the unopened product container at once.

By using this product user or buyer accepts the following Conditions Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions presence of other materials or the manner of use or application all of which are beyond the control of Control Solutions, Inc. To the extent consistent with applicable law all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Control Solutions, Inc. MAKES NO OTHER WARRANTIES EXPRESS OR IMPLIED OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein TO THE EXTENT CONSISTENT WITH APPLICABLE LAW CONTROL SOLUTIONS, INC. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT WARRANTY TORT NEGLIGENCE STRICT LIABILITY OR OTHERWISE SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT CONTROL SOLUTIONS, INC.'S ELECTION THE REPLACEMENT OF PRODUCT.

Impact is registered trademarks of Amvac Chemical Company

NOTES

NOTES

NOTES



NULLIFY A/P

ACTIVE INGREDIENT:

Glufosinate ammonium* 24.5%

Other Ingredients: 75.5%

Total: 100.0%

*Liquid soluble concentrate containing 2.34 pounds of active ingredient per U.S. gallon

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN THE EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not give any liquid to the person.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE AT 1-800-424-9300.	
NOTE TO PHYSICIAN: If this product is ingested endotracheal intubation and gastric lavage should be performed as soon as possible followed by charcoal and sodium sulfate administration.	

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label find someone to explain it to you in detail.)

See inside label booklet for additional PRECAUTIONARY STATEMENTS and DIRECTIONS FOR USE

EPA Reg. No. 42750-258-53883

(A) EPA Est. No. 88746-GA-01

(B) EPA Est. No. 53883-TX-002

First letter of lot# corresponds to EPA Est. # where product was produced

MANUFACTURED FOR: CONTROL SOLUTIONS, INC.
5903 Genoa-Red Bluff | Pasadena, TX 77507

EPA 080218