

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through the skin or inhaled. Skin sensitizer. Avoid contact with eyes, skin or clothing. Avoid inhaling vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**FIRST AID**

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

**NOTE TO USER:** Wear long sleeve shirt, long pants, gloves and goggles or face shield while mixing and applying this product. Keep unprotected persons and animals out of treatment area until spray has completely dried.

**ENVIRONMENTAL HAZARDS:**

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT apply directly to water, to areas where surface water is present, or intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378. Have product container or label with you when calling a poison control center or doctor or going for treatment.

For questions pertaining this product, please email us at support@southernag.com

**S A SOUTHERN  
5 0 AG®**

Contains  
**Daconil®**

**LIQUID  
ORNAMENTAL  
and  
VEGETABLE  
FLOWABLE FUNGICIDE**

**NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.**

**Active ingredient:**

Chlorothalonil (tetrachloroisophthalonitrile) ..... 12.50%

**Inert ingredients** ..... 87.50%

**Total** ..... 100.00%

Daconil® is a registered trademark of Syngenta Inc.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side panel and inside of label for additional precautionary information

**SHAKE WELL BEFORE USING.**

Read entire label and use as directed. Apply only to plants specified on this label. Do not apply to plants not listed on this label. Do not combine this product with pesticides, surfactants or fertilizers unless prior use has shown the combination compatible.

**Southern Agricultural Insecticides, Inc.**

Palmetto, Fla. 34220 Hendersonville, N.C. 28793 Boone, N.C. 28607

E.P.A. Reg. No. 829-287 J4 EPA Est No. 829-FL- 1

**Net contents liquid: 1 Quart (.946 liters)**

**STORAGE AND DISPOSAL:**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store pesticide in the original container in a locked storage area.


**PESTICIDE DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THE SPECIFIC DIRECTIONS FOR USE OF THIS PRODUCT ARE CONTINUED INSIDE OF LABEL.

FOLLOW DIRECTIONS ON SIDE OF LABEL TO OPEN.



0 51538 06823 9

## DIRECTIONS FOR USE (CONTINUED)

### GENERAL PRECAUTIONS AND RESTRICTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT allow people or pets to enter treated areas until sprays have dried or dusts have settled. DO NOT apply this product in a way that will contact other persons, or pets, either directly or through drift. DO NOT combine this product in the spray tank with other pesticides, surfactants, or fertilizers unless your prior use has shown the combination physically compatible, effective and non injurious under your conditions of use. The use of spray stickers or other spray adjuvants is not recommended with this product.

Before opening, slowly invert this container several times to assure uniform mixture. Slowly add the required amount of this product to the spray tank while filling. Agitate and continue to agitate while using.

For handy reference, 1 oz liquid = 2 Tablespoonfuls (TBS) = 6 Teaspoonfuls (tsp)

### APPLICATIONS INSTRUCTIONS

Use of this product on home lawns is prohibited. This product may not be added to paint by residential users/consumers.

## ORNAMENTAL PLANTS:

Apply this fungicide at the rate of 2 1/2 Tablespoonfuls (2 Tablespoonfuls for roses) per gallon of water to cover 350 sq. ft. Apply the spray to run-off when conditions are favorable for disease development. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable. During periods when conditions favor severe disease incidence, generally cloudy wet weather, apply at one week intervals. Apply to plants when foliage and flowers are dry or nearly dry.

Use of this product is recommended for the control of fungal diseases referred to by numbers in parentheses following each ornamental. Ornamentals listed on this label have been tested and found to tolerate applications of chlorothalonil (the active ingredient in this product) at the recommended rates. The user should test for possible phytotoxic responses, using recommended rates on ornamental plants on a small area prior to extensive use. Applications to bloom may damage flowers and/or fruits.

**FRUITS AND OTHER STRUCTURES WHICH ARE BORNE ON TREATED PLANTS MUST NOT BE EATEN** unless directions for use on the specific food plant are followed as given in the vegetable and fruit sections of this label. Do not feed treated plant parts to livestock. Do not apply more than 27 applications per growing season (year).

## ORNAMENTALS RECOMMENDED FOR TREATMENT WITH THIS FLOWABLE FUNGICIDE.

### BROADLEAF SHRUBS AND TREES

Andromeda(Pieris)(4), Ash(Fraxinus)(1), Aspen(1), Azalea(1,2,4), Buckeye(horsechestnut)(1), Cherry-Laurel(1), Crabapple(1,6,8), Dogwood(1), Eucalyptus(3), Euonymus(1), Firethorn(Pyracantha)(1), Flowering Almond(1,2), Flowering Cherry(1,2), Flowering Peach(1,2), Flowering Plum(1,2), Flowering Quince(1,2), Hawthorn(1,6), Holly (1), Lilac(5), Magnolia, Maple(1), Mountain Laurel(1), Oak(Red group only)(1,7), Oregon-Grape(Mahonia)(6), Photinia(1), Poplar(1), Privet(ligustrum)(1), Rhododendron(1,2,4), Sand Cherry(1,2), Sequoia(1), Spiraea(1), Sycamore,Planetree(1), Viburnum(5), Walnut(Juglans)(1)

### FLOWERING PLANTS AND BULBS

(Avoid applications during bloom period on plants where flower injury is unacceptable.)

Arabian Violet(2), Begonia(1), Camellia(2), Carnation(1,2), Chrysanthemum(1,2), Crocus(1), Daffodil(1), Daisy(1), Geranium(1,6), Gladiolus(1,2), Holly-hock(6), Hydrangea(Foliage only)(1,6), Iris(1,2), Lily(1), Marigold(1), Narcissus(1), Pansy(1), Petunia(1,4), Phlox(1), Poinsettia(1)\*Discontinue applications to poinsettia prior to bract formation; phytotoxicity is possible on the bracts, Rose(1), Statice(1), Tulip(1), Zinnia(1,5) Do not apply more than 30 applications per season (year).

### FOLIAGE PLANTS

Aglaoanema(1), Areca palm(1), Artemesia(1), Boston fern(1), Dumbcane(Diffenbachia)(1), Dracaena(1), Fatsia(Aralia)(1), Ficus(1), Florida Ruffle Fern(1), Leatherleaf Fern(1), Lipstick plant(1), Ming aralia(1), Oyster plant(Rhoeo)(1), Parlor palm, Chamaedorea(1), Peperomia(1), Philodendron(1,4), Prayer plant(Margantia)(1), Syngonium(1), Zebra plant(Aphelandra)(1).

## DISEASES CONTROLLED WITH THIS FLOWABLE FUNGICIDE

### (1) LEAFSPOTS/FOLIAR BLIGHTS

Actinopelle leafspot	Alternaria leafspot/leaf blight	Anthrachnose leaf blotch.spot	Anthrachnose(Discula)blight
Ascochyta blight	Bipolaris(Helminthosporium leaf spot)	Botrytis leafspot, leaf blight	Cephalosporium leaf spot
Cercospora leaf spot	Cercosporidium leafspot	Coryneum blight(shothole)	Corynespora leaf spot
Curvularia leaf spot	Cylindrosporium leafspot	Dactylaria leafspot	Didymellina leafspot
Drechslera leafspot	Fabraea(Entomosporium)leafspot	Fusarium leafspot	Gleosporium black leafspot
Ink spot(Drechslera)	Monilinia blossom blight, twig blight	Mycosphaerella ray blight	Myrothecium leafspot,
brown rot	Nematostoma leaf blight	Phyllosticta leaf spot	Rhizoctonia web blight
Ramularia leaf spot	Septoria leaf spot	Sphaeropsis leafspot	Stagonospora leaf scorch
Tan leafspot(Curvularia)	Volutella leaf blight		

### (2) FLOWER SPOTS/BLIGHTS

Botrytis flower spot, flower blight	Curvularia flower spot	Monilinia blossom blight
Uvulinia flower blight	Rhizopus blossom blight	Sclerotinia flower blight

### (3) CYLINDROCLADIUM STEM CANKER

### (4) PHYTOPHTHORA LEAF BLIGHT, dieback

### (5) POWDERY MILDEWS

Erysiphe cichoracearum	Microsphaera spp
------------------------	------------------

### (6) RUSTS

Gymnosporangium spp.	Puccinia spp.	Pucciniastrum hydrangeae
----------------------	---------------	--------------------------

### (7) TAPHRINA blister

### (8) SCAB (Venturia inaequalis)

## THE FOLLOWING ORNAMENTAL PLANT SPECIES WHICH HAVE BEEN TESTED WITH CHLOROTHALONIL AT RECOMMENDED RATES DID NOT EXHIBIT PHYTOTOXICITY.

<u>BOTANICAL NAME</u>	<u>COMMON NAME</u>	<u>BOTANICAL NAME</u>	<u>COMMON NAME</u>
Aechmea fasciata	Aechmea	Episcia cupreata	Flame Violet
Araucaria heterophylla	Norfolk Island Pine	Fittonia spp.	Silver-Nerve Plant
Asplenium nidus	Birdnest fern	Gerbera jamesonii	Gerbera Daisy
Bougainvillea spp.	Bougainvillea	Gynura sarmentosa	Purple Passion Vine
Caladium spp.	Caladium	Gypsophila paniculata	Baby's Breath
Calathea makoyana	Peacock plant	Hoya spp.	Wax Plant
Calistephus chinensis	Aster	Ilex cornuta	Chinese Holly
Carissa grandiflora	Natal plum	Ilex crenata	Japanese Holly
Clerodendron thomsonae	Bleeding heart	Impatiens spp.	Impatiens
Codiaeum spp.	Croton	Pilea cadierei	Aluminum Plant
Cordyline terminalis	TI Plant	Platycerium spp.	Staghorn fern
Crassula argentea	Jade Plant	Sansevieria trifasciata "Hahnii"	Birdnest Sansevieria
Cyrtomium falcatum	Holly Leaf Fern	Tolmeia menziesii	Piggy-back Plant
Dionaea muscipula	Venus Fly Trap	Yucca elephantipes	Spineless Yucca
Dizygotheca elegantissima	False Aralia	Zygocactus truncatus	Christmas cactus
Epipremnum aureum	Golden Pothos, Schindapsus		

**NOTE:** DO NOT apply this product to either green or variegated Pittosporum or to Schefflera, as multiple applications have been demonstrated to cause phytotoxic responses.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through the skin or inhaled. Skin sensitizer. Avoid contact with eyes, skin or clothing. Avoid inhaling vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**FIRST AID**

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

**NOTE TO USER:** Wear long sleeve shirt, long pants, gloves and goggles or face shield while mixing and applying this product. Keep unprotected persons and animals out of treatment area until spray has completely dried.

**ENVIRONMENTAL HAZARDS:**

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT apply directly to water, to areas where surface water is present, or intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas.

For questions pertaining this product, please email us at [support@southernag.com](mailto:support@southernag.com)

**S A SOUTHERN  
5 0 AG®**

Contains  
**Daconil®**

and

**LIQUID  
ORNAMENTAL  
VEGETABLE  
FLOWABLE FUNGICIDE**

**NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.**

**Active ingredient:**

Chlorothalonil (tetrachloroisophthalonitrile) ..... 12.50%

Inert ingredients ..... 87.50%

**Total** ..... 100.00%

Daconil® is a registered trademark of Syngenta Inc.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side panel and inside of label for additional precautionary information

**SHAKE WELL BEFORE USING.**

Read entire label and use as directed. Apply only to plants specified on this label. Do not apply to plants not listed on this label. Do not combine this product with pesticides, surfactants or fertilizers unless prior use has shown the combination compatible.

**Southern Agricultural Insecticides, Inc.**

Palmetto, Fla. 34220 Hendersonville, N.C. 28793 Boone, N.C. 28607  
E.P.A. Reg. No. 829-287 J3 EPA Est No. 829-FL- 1

**Net contents liquid: 1 Quart (.946 liters)**

**STORAGE AND DISPOSAL:**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store pesticide in the original container in a locked storage area.

**PESTICIDE DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378. Have product container or label with you when calling a poison control center or doctor or going for treatment

.For questions pertaining this product, please email us at [support@southernag.com](mailto:support@southernag.com)

## ORNAMENTAL CONIFERS:

Apply this fungicide at the rate given below per gallon of water to cover 350 sq. ft.

Apply to obtain uniform coverage of tree canopy. Do not allow livestock to graze treated areas.

Rates are given in Tablespoonfuls of product per gallon of water.

Do not make more than 6 applications per year for rates up to 4 1/2 TBS. Do not make more than 4 applications per year at rates above 4 1/2 TBS.

DISEASES	TBS/GAL	APPLICATION DIRECTIONS
Swiss needlecast	7 1/2 TBS.	Make first application in
Scleroderris canker(pines)	2 1/2 TBS to 4 1/2 TBS	Spring when new shoot growth is 1/2 to 2 inches long. Repeat applications may be made after 21 days. Make additional applications at 3 to 4 week intervals
Sirococcus tip blight	3 1/2 TBS	until conditions no longer favor disease development.

Cyclaneusma and Lophodermium needlecasts	7 1/2 TBS	Apply in early Spring prior to bud break. Repeat applications at 6 to 8 week intervals, until spore release ceases in late Fall. Apply monthly during periods of frequent rainfall. During drought periods applications may be suspended, then resumed upon next occurrence of needle wetness.
--	-----------	--

Rhabdocline needlecast	2 1/2 TBS to 4 1/2 TBS	Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development.
------------------------	------------------------	---

Botrytis seedling blight Phoma twig blight	2 1/2 TBS	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development.
---	-----------	--

## TREE AND ORCHARD CROPS:

Dosages are given in Tablespoonfuls (TBS) per gallon of water. Spray to point of run-off. Obtain complete coverage of the tree canopy.

For stone fruit do not use more than 2.5 gallons of spray per 350 sq. ft. Do not exceed 5 applications per season.

For papaya do not use more than 1 gallon of spray per 350 sq. ft. Do not exceed 3 applications per season. **DO NOT** apply after shuck split and before harvest.

If additional fungicide applications are needed before harvest, apply an alternate fungicide labeled for that use.

CROP	DISEASES	TBS/GAL	APPLICATION DIRECTIONS
STONE FRUIT (Peach, Nectarine, Apricot, Cherry, Plum, Prune)	Leaf curl, Coryneum blight (shothole)	2TBS	Apply at leaf fall in Autumn. Do not retreat within 10 days. When conditions favor disease make one or two additional applications in mid and late Winter before budswell. Where Coryneum blight (shothole) occurs also apply at budbreak and shuck split.
	Brown rot blossom blight Lacy (russet) scab (plum/prune)	2TBS	Apply once at popcorn (pink, red, or early white bud) and a second application at full bloom. If weather conditions favor disease development make an additional application at petal fall. Do not retreat within 10 days.
	Cherry leaf spot; Peach, Nectarine, Apricot scab	2 TBS	In addition to the bloom applications make one application at shuck split. For control of Cherry leaf spot after harvest, make one application 7 days after fruit has been removed. Where trees have a history of high leaf spot incidence, make a second application 10-14 days later.
Papaya	Alternaria, Fruit rot Anthracnose Stem end rot	3 TBS to 4 TBS	Begin treatment when conditions favor disease development. Do not retreat within 14 days.

## VEGETABLE DISEASES:

Rates are given in Tablespoonfuls (TBS) per gallon of water to cover 350 square feet of planting. Spray to obtain thorough coverage but do not exceed 1 gallon of spray mixture per 350 square feet or residue tolerances may be exceeded. The number shown in parentheses following crop name indicates the interval between last application and harvest. When sprays for disease control must be made at intervals closer to harvest than indicated use an alternate fungicide registered for that use. **DO NOT** graze treated areas or feed treated plant parts to livestock.

CROP	DISEASES	TBS/GAL	APPLICATION DIRECTIONS
Bean (snap)(7)	Rust Botrytis blight (gray mold)	2 TBS to 4 TBS 4 TBS	Begin applications during early bloom stage or when disease first threatens. Do not retreat within 7 days. Repeat weekly or as necessary to maintain control. Do not exceed 4 applications per season.
Beans (dry)(402) Navy, Pinto,	Rust Anthracnose	2 TBS	Begin applications during early bloom stage. Do not retreat within 7 days. For use only on beans to be harvested dry with pods removed. Do not exceed 5 applications per season.
Kidney, Lima, Blackeye	Downy Mildew Cercospora leaf spot (blackeye only)		
Cabbage Chinese Cabbage (7) (tight headed varieties only) Cauliflower Broccoli Chinese Broccoli (7) Brussels sprouts	Alternaria leaf spot Downy mildew	2 1/2TBS	Begin applications when transplants are set in field, or shortly after emergence for field seeded crop, or when conditions favor disease development. Do not retreat within 7 days. Repeat at 7-10 day intervals, or as necessary to maintain control. Do not exceed 8 applications per season.
Carrot (7)	Cercospora (early blight) Alternaria (late blight)	2 1/2 TBS	Start applications when disease threatens. Do not retreat within 7 days. Repeat applications at 7-10 day intervals or as long as necessary to maintain control. Do not exceed 11 applications per season.
Celery(7)	Cercospora (early blight) Septoria (late blight) Basal stalk rot (Rhizoctonia solani)	3 TBS to 4TBS	Weekly applications when transplants are set in field. Do not exceed 8 applications per season. Do not retreat within 7 days.
Corn(sweet)(14)	Helminthosporium leaf blights, Rust	1 TBS to 2 1/2TBS	Begin applications when conditions favor disease development. Do not retreat within 7 days. Repeat at 7 to 10 day intervals or as necessary to maintain control. Do not exceed 6 applications per season.

**DO NOT APPLY TO SWEET CORN TO BE PROCESSED. DO NOT ALLOW LIVESTOCK TO GRAZE IN TREATED FIELDS. DO NOT ENSILE TREATED CORN OR USE AS LIVESTOCK FORAGE.**

## VEGETABLE DISEASES: (continued)

CUCURBITS:	Anthraxnose	2 TBS	Begin applications when plants are in first true leaf stage, or when conditions favor disease development. Do not retreat within 7 days. Do not exceed 7 applications per season.
Cucumber	Downy Mildew	to 2 1/2 TBS	
Cantaloupe	Target spot	3 TBS to 4 TBS	Under severe conditions shorten spray intervals.
Muskmelon	Cercospera		
Honeydew	Leaf spot		
Watermelon	Gummy stem blight		
Squash	(black rot)		
	Scab		
Pumpkin	Powdery Mildew(Sphaerotheca only)		
Onion (dry bulb)(7)	Botrytis leaf blight/blast	1 1/2 TBS to 2 1/2 TBS	Do not retreat within 10 days. Prior to infection shorten spray interval to 7 days when <u>disease is present</u> . Do not exceed 11 applications per season.
	Botrytis neck rot (suppression)		
	Purple blotch		
	Neck rot in storage (suppression)		A minimum of 3 applications prior to one week before lifting.
Onion (green bunching)(14)*, Leak (14)*, Shallot(14)*, Garlic (7)	Botrytis leaf blight/blast Purple blotch Downey mildew (suppression)	2 TBS to 4 TBS	Begin weekly applications prior to favorable infection periods and continue for as long as conditions favor disease. Do not retreat within 7 days. DO NOT apply more than 3 times per season to green bunching onions, leeks, or shallots. For Garlic do not exceed 6 applications per season.
Parsnip(10)	Alternaria leaf	2 1/2 TBS	Make first application at first sign of disease or when conditions favor disease infection. Do not retreat within 7 days. DO NOT apply more than 4 times per season.
Peanut(14)	Early leaf spot (Cercospera) Late leaf spot (Cercosporidium) Rust, Web blotch.	2 TBS	Apply when leaf wetness first occurs or 30-40 days after planting. Do not retreat within 10 days. Do not exceed 8 applications per season.
Potato (7)	Late blight	2 TBS	Apply as a banded treatment directed over the foliage, beginning when vines are first exposed and leaf wetness occurs. Do not retreat within 5 days. As vines close apply as a complete cover spray at 7-10 day intervals. Use the 7 day interval when disease conditions worsen, being sure to apply 1 gallon for each 350 square feet. Do not exceed 10 applications per season.

Tomato	Early blight, Late blight, Gray Leaf Spot, Gray Leaf mold, Septoria leaf spot, Target spot, Anthracnose, Alternaria fruit rot, Botrytis gray mold, Rhizoctonia fruit rot	3 TBS to 4 TBS	Do not retreat within 7 days. Begin applications when dew or rain occurs and disease threatens. Use shorter interval as disease conditions worsen. Do not exceed 7 applications per season.
	to aid in the control of bacterial foliar and fruit spots		add 4 teaspoonfuls of SA-50 Neutral Copper Fungicide or add 2 Tablespoonfuls of SA-50 Liquid Copper Fungicide per gallon of spray mixture.

## WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with its directions under normal conditions of use and buyer assumes the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. In no event shall the seller's breach of warranty exceed the purchase price of the material as to which a claim is made.

