

BASAGRAN® T/O HERBICIDE

We Don't Make The Turf.
We Make It Better.™

 **BASF**
The Chemical Company



Better Postemergent Control

Basagran® T/O herbicide offers reliable, cost-effective postemergent control of tough broadleaf weeds, yellow nutsedge, annual sedges and more. It can also provide weed control in ornamentals. When used as directed, **Basagran T/O:**

- Provides a cost-effective nutsedge control program.
- Effectively eliminates underground tubers.
- Delivers effective postemergence activity, which allows flexibly timed weed control.
- Can be used in cool- and warm-season turfgrass and ornamentals.
- Can be easily tank-mixed with other herbicides to affordably increase their spectrum of control.

Weeds Controlled

In established cool- and warm-season turfgrasses, ornamentals and roadsides, **Basagran T/O** controls:

- | | |
|-------------------|----------------|
| • Yellow Nutsedge | • Wild Mustard |
| • Annual Sedges | • Spurweed |
| • Common Purslane | • Ragweed |
| • Lawn Burweed | • Galinsoga |
| • Musk Thistle | • Dayflower |
| • Canada Thistle | |

Use Sites:

- Landscape or Grounds Maintenance
 - Noncropland Areas
 - Christmas Trees and Other Forest Tree Nurseries
 - Non-Bearing Fruit, Nut and Vines
 - Golf Courses
 - Residential and Commercial Lawns
 - Sod Farms
 - Athletic Fields
 - Field and Container Nurseries
- Do not apply to putting greens.*

Formulation: 4 SL
(Soluble Liquid)

Packaging: 2.5 gallons

Active Ingredient: Sodium Bentazon

Chemical Family: Benzothiadiazole

Mode of Action: Photosynthetic Inhibitor complex II in chloroplast. The plant's ability to make food is disrupted and the weed dies.

Rates: 0.75 oz. / 1000 sq. ft. (2 pints / A) in 1 to 2 gallons of water / 1000 sq. ft. (40 to 80 gallons / A).

Control Symptoms: Chlorosis begins 3 to 5 days following application followed by necrosis and desiccation of the plant.

Basagran[®] Herbicide
T/O

Keys to Success

Basagran T/O needs to be combined with a crop oil concentrate (COC) at 0.75 oz. / 1000 sq. ft. (2 pints / A) for performance on targeted species. Applications to ornamentals should be directed sprays around the base of the plant. Do not mow 3 days before or after application. Repeat application in 10 to 14 days to improve long-term results.

Suggested Tank Mixes

Atrazine, MSMA, 2,4-D, 2,4-DP and MCPP are compatible with most of the commonly used broadleaf weed herbicides in turf. (Do not apply oil concentrate with 2,4-D, 2,4-DP or MCPP tank mixes.) Consult all product label restrictions.

Always read and follow label directions.

Basagran is a registered trademark of BASF. We Don't Make The Turf. We Make It Better. is a trademark of BASF.

©2006 BASF Corporation. All rights reserved. APN 05-14-002-0108J

Basagran[®] Herbicide
T/O