



Syngenta Bravo Weather Stik®



Product Overview

Bravo Weather Stik® fungicide is a world-leading broad-spectrum foliar protectant fungicide labeled for use on vegetables. It has a patented Weather Stik® formulation that sticks and stays on the plant surface longer than any other preventive fungicide available to provide superior broad-spectrum disease control. Bravo Weather Stik stays on to fight disease, even after heavy rains or irrigation. Bravo® fungicide is also available in granular formulation as Bravo Ultrex® and a liquid ZN premix as Bravo ZN.

Key Benefits

- Controls most major diseases on a variety of vegetables
- Patented Weather Stik trait provides superior rainfastness
- Provides long-lasting residual control
- Effective rotational product for disease resistance management because of its multi-site mode of action that greatly reduces the likelihood of fungi developing resistance

Application Details

- Apply by ground, air or chemigation, but be sure to consult the product label for the approved application method per crop.
- Use Bravo Weather Stik in sufficient water to obtain thorough coverage of foliage.
- Begin applications prior to conditions that favor disease development.
- Repeat application as necessary while maintaining a seven-day interval between applications.
- Bravo Weather Stik is an extreme eye irritant. Follow label instructions regarding re-entry into a treated field.

Bravo Weather Stik LABEL AT A GLANCE

Vegetables*			
Crop	Diseases (Pathogen)	Use Rate pts. product/A (lbs. a.i./A)	Preharvest Interval (PHI)
Asparagus	Cercospora Blight (<i>C. asparagi</i>), Purple Spot (<i>Pleospora herbarum</i>), Rust (<i>Puccinia asparagi</i>)	2.0 - 4.0 (1.5 - 3.0)	190 ¹
Bean (snap)	Rust (<i>Uromyces appendiculatus</i>)	1.375 - 3.0 (1.0 - 2.25)	7
	Botrytis Blight (<i>B. cinerea</i>)	3.0 (2.25)	
Broccoli Cabbage Cauliflower	Alternaria Leaf Spot (<i>Alternaria</i> spp.), Downy Mildew (<i>Peronospora parasitica</i>)	1.5 (1.125)	7
	Ring Spot (California only)	2.0 (1.5)	
Carrot	Alternaria Leaf Blight (<i>A. dauci</i>), Cercospora Leaf Spot (<i>C. carotae</i>)	1.5 - 2.0 (1.125 - 1.5)	0



Bravo Weather Stik LABEL AT A GLANCE

Vegetables* continued

Crop	Diseases (Pathogen)	Use Rate pts. product/A (lbs. a.i./A)	Preharvest Interval (PHI)
Celery	Basal Stalk Rot (<i>Rhizoctonia solani</i>), Early Blight (<i>Cercospora apii</i>), Late Blight (<i>Septoria apicola</i>)	2.0 - 3.0 (1.5 - 2.25)	7
	Suppression: (Seven-day schedule): Pink Rot (<i>Sclerotinia sclerotiorum</i>)	3.0 (2.25)	
	Early Blight (<i>Cercospora apii</i>), Late Blight (<i>Septoria apicola</i>)	1.5 - 2.0 (1.125 - 1.5) /100 gals.	
Corn (sweet and grown for seed)	Helminthosporium Leaf Blights Rust (<i>Puccinia</i> spp.)	0.75 - 2.0 (0.6 - 1.5)	14
Cucurbits	Anthrachnose (<i>Colletotrichum</i> spp.), Downy Mildew (<i>Pseudoperonospora cubensis</i>), Target Spot (<i>Corynespora cassiicola</i>)	1.5 - 2.0 (1.125 - 1.5)	0
	Alternaria Leaf Blight (<i>A. cucumerina</i>), Alternaria Leaf Spot (<i>A. alternata</i>), Cercospora Leaf Spot (<i>C. citrullina</i>), Gummy Stem Blight/Vine Decline (<i>Didymella bryoniae</i>), Powdery Mildew (<i>Sphaerotheca</i> only), Scab (<i>Cladosporium cucumerinum</i>)	2.0 - 3.0 (1.5 - 2.25)	
Onions (dry bulb) Garlic	Botrytis Leaf Blight (<i>Botrytis</i> spp.), Purple Blotch (<i>Alternaria porri</i>), Suppression: Botrytis Neck Rot, Downy Mildew (<i>Peronospora destructor</i>)	1.0 - 3.0 (0.75 - 2.25)	7
Onion (green bunching) Leek Shallots Onion and Garlic (grown for seed)	Botrytis Leaf Blight (<i>Botrytis</i> spp.), Purple Blotch (<i>Alternaria porri</i>), Suppression: Downy Mildew (<i>Peronospora destructor</i>)	1.5 - 3.0 (1.125 - 2.25)	14 ²
Tomato	FOLIAGE: Early Blight (<i>Alternaria solani</i>), Gray Leaf Mold (<i>Fluvia fluva</i> ; <i>Cladosporium</i>), Gray Leaf Spot (<i>Stemphyllium botryosum</i>), Late Blight (<i>Phytophthora infestans</i>), Septoria Leaf Spot (<i>S. lycopersici</i>), Target Spot (<i>Corynespora cassiicola</i>)	1.375 - 2.0 (1.0 - 1.5)	0
	FRUIT: Alternaria Fruit Rot (<i>A. alternata</i>), Anthracnose (<i>Colletotrichum</i> spp.), Botrytis Gray Mold (<i>B. cinerea</i>), Late Blight Fruit Rot (<i>P. infestans</i>), Rhizoctonia Fruit Rot (<i>R. solani</i>)	2.0 - 2.75 (1.5 - 2.1)	

* Consult the product label for complete application directions.

¹ Do not apply within 190 days (120 days in California and Arizona) of the harvest of spears in the following season.

² Do not apply within seven days of harvest on garlic.



For more information, visit www.farmassist.com, www.syngentacropprotection.com or call the Syngenta Customer Center at 1-866-SYNGENT(A) (796-4368).

©2009 Syngenta Crop Protection, Inc., P.O. Box 18300, Greensboro, NC 27419.

Important: Always read and follow label instructions before buying or using these products.

Syngenta Crop Protection, Inc. warrants that its products conform to the chemical description set forth on the products' labels. NO OTHER WARRANTIES, WHETHER EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF MERCHANTABILITY AND OF FITNESS FOR A PARTICULAR PURPOSE, SHALL APPLY TO SYNGENTA PRODUCTS. Syngenta Crop Protection, Inc. neither assumes nor authorizes any representative or other person to assume for it any obligation or liability other than such as is expressly set forth herein. UNDER NO CIRCUMSTANCES SHALL SYNGENTA CROP PROTECTION, INC. BE LIABLE FOR INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF ITS PRODUCTS. No statements or recommendations contained herein are to be construed as inducements to infringe any relevant patent now or hereafter in existence.

Bravo®, Bravo Ultrax®, Bravo Weather Stik®, Weather Stik® and the Syngenta logo are registered trademarks of a Syngenta Group Company.

G&S 409.60225 (05/09)

SCP 123-00039-A