

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)

GORDON'S® TRIMEC® CRABGRASS PLUS LAWN WEED KILLER CONCENTRATE

*Also Kills Dandelion & Clover
Plus 200 Other Listed Weeds*

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	6.42%
Quinclorac	2.13%
Dicamba, dimethylamine salt	0.60%
INERT INGREDIENTS:	90.85%
TOTAL 100.00%	

THIS PRODUCT CONTAINS:
 0.456 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%
 0.182 lb. 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%
 0.043 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%
 Isomer Specific By AOAC Methods.

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING

1. WHAT IT DOES: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See below for complete list.)

2. HOW IT WORKS: Trimec® Crabgrass Plus Lawn Weed Killer Concentrate contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

3. WHERE TO USE:

Trimec Crabgrass Plus Lawn Weed Killer Concentrate can be applied to residential lawns with the following turfgrass species:

Lawns To Be Treated:
Cool-Season Turfgrass
Kentucky bluegrass, Ryegrass (perennial and annual), Tall fescue, Mixtures of cool season grasses containing fine fescues (red, hard and chewings).
Warm-Season Turfgrass
Bermudagrass ¹ (common and hybrid), Zoysiagrass, Buffalograss.
¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

DO NOT apply Trimec Crabgrass Plus Lawn Weed Killer Concentrate to the following:
Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens including vegetables, fruit trees, vines and berries, Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore paspalum, St. Augustinegrass, Turfgrass species that are not listed on this label.

4. GRASS AND BROADLEAF WEEDS CONTROLLED:

A partial list is presented below:

	GRASS WEEDS	
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf
	BROADLEAF WEEDS	
Annual yellow sweetclover	Buckhorn plantain	Cocklebur
Aster	Bulbous buttercup	Common mullein
Austrian fieldcress	Bull thistle	Creeping Jenny
Bedstraw	Bullnettle	(Creeping Charlie)
Beggarticks	Burclover	Cudweed
Betony, Florida	Burdock	Curly dock
Bindweed, field	Burning nettle	Daisy, English
Bird vetch	Burweed	Daisy fleabane
Bitter wintercress	Buttercup	Daisy, oxeley
Bittercress, hairy	Buttonweed	Dandelion
Bitterweed	Canada thistle	Dichondra
Black-eyed Susan	Carolina geranium	Dogbane
Black medic	Carpetweed	Dogfennel
Black mustard	Catchweed bedstraw	Dollarweed
Blackseed plantain	Catnip	Elderberry
Blessed thistle	Catsear	False dandelion
Bloodflower milkweed	Chickweed, common	Falseflax
Blue lettuce	Chickweed, mouseear	False sunflower
Blue vervain	Chicory	Fiddleneck
Bracted plantain	Cinquefoil	Florida pusley
Brassbuttons	Clover, crimson	Frenchweed
Bristly oxtongue	Clover, hop	Galinsoga
Broadleaf dock	Clover, red	Goathead
Broadleaf plantain	Clover, strawberry	Goldenrod
Broomweed	Clover, sweet	Ground ivy
Buckhorn	Clover, white	Gumweed
	Cockle	

(continued on next page)

BROADLEAF WEEDS (cont.)

Hairy fleabane	Plantain	Tansy mustard
Hawkweed	Poison ivy	Tanweed
Healall	Poison oak	Thistle
Heartleaf drymary	Pokeweed	Trailing crownvetch
Heathaster	Poorjoe	Tumble mustard
Hedge bindweed	Prairie sunflower	Tumble pigweed
Hedge mustard	Prickly lettuce	Velvetleaf
Hemp	Prickly sida	Venice mallow
Henbit	Prostrate knotweed	Virginia buttonweed
Hoary cress	Prostrate pigweed	Virginia creeper
Hoary plantain	Prostrate spurge	Virginia pepperweed
Hoary vervain	Prostrate vervain	Water pennywort
Horsenettle	Puncturevine	Wavyleaf bullthistle
Jimsonweed	Purslane, common	Western clematis
Knawel	Ragweed	Western salsify
Knotweed	Red sorrel	White mustard
Kochia	Redroot pigweed	Wild aster
Lambsquarters	Redstem filaree	Wild buckwheat
Lespedeza	Rough cinquefoil	Wild carrot
Mallow	Rough fleabane	Wild four-o'clock
Matchweed	Russian pigweed	Wild garlic
Mexicanweed	Russian thistle	Wild geranium
Milk vetch	Scarlet pimpernel	Wild lettuce
Morningglory	Scotch thistle	Wild marigold
Mouseear hawkweed	Sheep sorrel	Wild mustard
Mugwort	Shepherdspurse	Wild onion
Musk thistle	Slender plantain	Wild parsnip
Mustard	Smallflower galinsoga	Wild radish
Narrowleaf plantain	Smooth dock	Wild rape
Narrowleaf vetch	Smooth pigweed	Wild strawberry
Nettle	Sorrel	Wild sweet potato
Orange hawkweed	Sowthistle	Wild vetch
Oriental cocklebur	Spanishneedles	Wild violet
Oxalis	Speedwell	Woodsorrel
Parsley-piert	Spiny amaranth	Woolly croton
Parsnip	Spiny cocklebur	Woolly morningglory
Pearlwort	Spiny sowthistle	Woolly plantain
Pennycress	Spotted catsear	Wormseed
Pennywort	Spotted spurge	Yarrow
Peppergrass	Spurweed	Yellow rocket
Pepperweed	Stinging nettle	Yellowflower pepperweed
Pigweed	Strawberry, India mock	and other broadleaf weeds
Pineywoods bedstraw	Tall nettle	
Plains coreopsis (tickseed)	Tall vervain	
	Tansy ragwort	

5. WHEN TO USE:

Apply this product when weeds are small and actively growing.

Weed species	Season	Timing	Maximum number of applications ¹	Application interval, days	Comments
Crabgrass (large and smooth):	Spring	From crabgrass emergence until the crabgrass is 3 to 4 inches in height.	2	14	Re-treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height.	2	14	Re-treat at the same rate at 14 days later for dense populations.
Broadleaf signalgrass	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	14	Re-treat at the same rate at 14 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing.	2	—	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing.	2	—	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

¹The maximum application rate is 6.4 fl. oz. of product per 1,000 sq. ft. per application (1.0 lb. 2,4-D acid equivalent per acre per application).
The maximum number of broadcast applications is limited to 2 per year. Do not apply more than 2 applications per year. The maximum seasonal rate is 12.8 fl. oz. of product per 1,000 sq. ft. (2.0 lbs. 2,4-D acid equivalent per acre).

6. HOW MUCH TO USE:

A. ENTIRE LAWN TREATMENTS WITH DIAL TYPE HOSE-END SPRAYERS

Three Easy Steps To Mix And Apply:

1. FILL

- Determine the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- The application rate for this product is 6.4 fl. oz. of product per 1,000 sq. ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area, or add 6.4 fl. oz. of product for every 1,000 sq. ft. of lawn area. Over-application or rates above those recommended on this label can cause turf injury.
- Do not add water to the sprayer jar.

2. SET THE DIAL

- Set the dial to the rate per gallon and keep the dial set to this setting while spraying.
- The dial setting or rate per gallon for this product is 2.5 fl. oz. of product per gallon of water.

Types of hose-end sprayers	Dial setting
Ortho® Dial'N Spray® Hose-End Sprayer	2.5 fl. oz.
Hudson® Hose End Liquid Concentrate Sprayer	2.5 fl. oz.
Gilmour® Dial-A-Mix® Lawn and Garden Sprayer	15 teaspoons or 5 Tablespoons

3. SPRAY

- Apply evenly over the lawn.

B. HAND-OPERATED SPRAYERS (backpack, knapsack, tank-type or pump-up):

- Apply to actively growing grass and broadleaf weeds during the early growth stages.
- Measure and mix the proper amount of product with water as described below. Mix 2.5 fl. oz. of product with 1 gallon of water to treat 400 sq. ft. of lawn. See below for additional spray preparations
- Spray evenly over the lawn. Spray coverage should be uniform and complete. Over-application or rates above those recommended on this label can cause turf injury.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Approximate Treated Area, Square Feet
2.5 fl. oz.	1 gallon	400 sq. ft.
3.75 fl. oz.	1.5 gallons	600 sq. ft.
5.0 fl. oz.	2 gallons	800 sq. ft.
7.5 fl. oz.	3 gallons	1,200 sq. ft.

Equal measures: 2.5 fl. oz. = 5 Tablespoons.
1 cup = 8 fl. oz. = 1/2 pint.

7. CULTURAL TIPS AND NOTES:

When to Apply:	For established lawns For newly seeded areas For newly sodded, sprigged, or plugged areas	<ul style="list-style-type: none"> • Apply when daytime temperatures are below 90°F. • Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing. • Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
Seeding:		<ul style="list-style-type: none"> • The treated area can be seeded at 4 weeks after the application of this product.
Irrigation:	Non-stressed turfgrass Stress from high temperatures, drought, insects or diseases	<ul style="list-style-type: none"> • Adequate soil moisture and favorable growing conditions enhance the performance of this product. • Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.
Mowing:		<ul style="list-style-type: none"> • Delay mowing 2 days before and until 2 days after the application of this product. • Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
Nontarget plants:		<ul style="list-style-type: none"> • Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. • Do not spray exposed roots of ornamentals and trees.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children or pets. Avoid contamination of food or feedstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call you local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

761/8-2009 AP081209
EPA REG. NO. 2217-896



MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com

SPECIMEN
LABEL